




Resilience Among Victims of Child Maltreatment

A Bibliography

July 2026

**Championing and Strengthening the
Global Response to Child Abuse**

nationalcac.org | 256-533-KIDS(5437) | 210 Pratt Avenue NE, Huntsville, AL 35801

©2026 National Children's Advocacy Center. All rights reserved.

© 2013, 2016, 2018, 2020, 2023, 2026. National Children's Advocacy Center.
All rights reserved.

Preferred citation: National Children's Advocacy Center. (2026). *Resilience among victims of child maltreatment: A bibliography*.

This project was supported by a grant awarded by the Office of Juvenile Justice and Delinquency Prevention, Office of Justice Programs, U.S. Department of Justice. Points of view or opinions in this document are those of the author and do not necessarily represent the official position or policies of the U.S. Department of Justice.

Scope

This bibliography provides research literature covering a wide variety of topics related to the resilience of child maltreatment victims.

Organization

Publications include articles, book chapters, reports, and research briefs and are listed in date descending order. Links are provided to full text publications when possible. However, this collection may not be complete. More information can be obtained in CALiO™, the Child Abuse Library Online.

Disclaimer

This bibliography was prepared by the Digital Information Librarians of the National Children's Advocacy Center (NCAC) for the purpose of research and education, and for the convenience of our readers. The NCAC is not responsible for the availability or content of cited resources. The NCAC does not endorse, warrant or guarantee the information, products, or services described or offered by the authors or organizations whose publications are cited in this bibliography. The NCAC does not warrant or assume any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed in documents cited here. Points of view presented in cited resources are those of the authors, and do not necessarily coincide with those of the NCAC.

Resilience Among Victims of Child Maltreatment

A Bibliography

Antunes, C., Camilo, C., Ferreira, C., Cunha, C., & Magalhães, E. (2026). [A strengths-based approach to resilience in child sexual abuse: A meta-analysis](#). *International Journal on Child Maltreatment: Research, Policy and Practice*, 9(2), 367–390.
DOI:10.1007/s42448-026-00255-5

This meta-analysis aims to uncover which protective factors reveal the greatest effect sizes on resilience among CSA survivors (i.e., symptoms, well-being, and competence) based on the Resilience Portfolio Model. This perspective shifts the focus from deficit-based models to a strengths-based approach that focuses on CSA survivors' resources & assets. Following the PRISMA guidelines, a total of 50 reports were included in the meta-analysis, involving 12,345 child & adult participants from different contexts (community, clinical, and forensic) and 335 effect sizes. Our findings revealed that while meaning-making strengths (e.g., spirituality) were associated with both well-being & symptoms, regulatory strengths (e.g., emotional regulation, self-control) & supportive relationships (e.g., caregiver emotional support) were associated with symptoms but not with well-being, and environmental strengths (e.g., extracurricular activities) were associated with well-being but not symptoms. Coping was not significantly associated with symptoms or well-being. Clinical interventions based on narrative, mindfulness, and trauma-focused therapy could be particularly useful for fostering psychological health in CSA victims.

Dominguez, D., Pérez-Testor, C., Benedico-Peydró, P., Casarramona, A., & Aznar-Martínez, B. (2025). [Assessing child maltreatment and its relationships with personality, resilience and attachment in adulthood](#). *Children & Society*, 39(4), 845–853.
DOI:10.1111/chso.12954

The main objective of this study is to examine the psychological effects of childhood abuse in adulthood, focusing on personality traits, resilience and attachment style within

a large sample specifically centred on the Spanish context. A total of 700 adults participated, of whom 24.4% reported experiencing some form of childhood abuse. The results indicate that individuals who reported experiencing childhood abuse exhibit lower levels of responsibility, agreeableness, extraversion and resilience, as well as a greater tendency to develop an insecure attachment style compared to those without a history of abuse. Furthermore, findings suggest that both attachment style and specific personality traits (extraversion, agreeableness and responsibility) partially mediate the relationship between childhood maltreatment and resilience. These results highlight the critical need for a comprehensive approach to prevent childhood abuse and neglect, emphasising the importance of early interventions to foster adaptive personality development, resilience and secure attachment in adulthood.

Fares-Otero, N. E., Carranza-Neira, J., Womersley, J. S., Stegemann, A., Schalinski, I., Vieta, E., Spies, G., & Seedat, S. (2025). [Child maltreatment and resilience in adulthood: A systematic review and meta-analysis](#). *Psychological Medicine*, 55, e163.
DOI:10.1017/S0033291725001205

We conducted a systematic review and meta-analysis to quantify associations between overall and subtypes of CM, global/trait resilience, and five resilience domains (coping, self-esteem, emotion regulation, self-efficacy, and well-being) in adults, and to examine moderators and mediators of these associations. A systematic search was undertaken on 12 June 2024 to identify published peer-reviewed articles in five databases (PROSPERO-CRD42023394120). Of 15,262 records, 203 studies were included, comprising 145,317 adults ($M_{age} = 29.62$ years; 34.96% males); 183 studies and 557 effect sizes were pooled in random-effect meta-analyses. Overall CM and its subtypes were negatively associated with global/trait resilience and its domains ($r = -0.081$ to -0.330). Emotional abuse/neglect showed the largest magnitude of effect ($r = -0.213$ to -0.321). There was no meta-analytic evidence for an association between sexual abuse and coping, and physical abuse/neglect and self-esteem. Meta-regressions identified age, sample size,

and study quality as moderators. Subgroup analyses found that associations between emotional abuse and emotion regulation were stronger, while associations between emotional abuse and self-esteem were weaker, in western *versus* non-western countries. No differences were found in associations between CM and resilience in clinical *versus* non-clinical samples. Narrative synthesis identified several mediators. Associations were of small magnitude and there were a limited number of studies, especially studies assessing CM subtypes, such as physical neglect, bullying, or domestic violence, and resilience domains, such as coping or self-efficacy, in males, and clinical samples. CM exposure negatively impacts resilience in adults, an effect observed across multiple maltreatment types and resilience domains. Interventions focused on resilience in adults with CM histories are needed to improve health and psychosocial outcomes.

van der Walt, C., & Mawila-Chauke, D. (2025). ["I had to be brave": Unveiling the remarkable resilience of children who experienced sexual abuse](#). *Journal of Child & Adolescent Trauma*, 18(3), 621–635. DOI:10.1007/s40653-025-00714-0

Sexual abuse is a global problem with profound consequences for the well-being of children. It often results in mental health disorders such as depression, post-traumatic stress disorder (PTSD) & anxiety. It can lead to social withdrawal, substance use & abuse, aggression and suicide. Despite the adversities faced by children who have experienced sexual abuse, some are resilient and do not succumb to the subsequent consequences. Therefore, this South African study aimed to explore the protective factors that enable the resilience of children who have been sexually abused. In line with the qualitative research, a phenomenological research design was used. Six female children in 2 children's homes were purposefully chosen for this study. Ungar's theory of resilience was the theoretical framework that underpinned this study. Data was collected through semi-structured interviews and analysed using Braun and Clarke's six-step thematic approach. The findings, based on participants' verbatim responses, identified individual, relational, and

contextual factors as essential social-ecological resources that support resilience in children who have experienced sexual abuse. Individual factors included qualities such as a capacity to help others, personal interests, bravery, and a forward-looking vision. Relational factors involved supportive social connections and caregiver support, while contextual factors included role models, religion & spirituality, and support from teachers. These findings highlight the importance of stakeholders in children's social ecology in developing and implementing supportive measures.

Panagou, C., & Macbeth, A. (2024). [Trajectories of risk and resilience: The role of empathy and perceived social support in the context of early adversity](https://doi.org/10.1016/j.chiabu.2024.106811). *Child Abuse & Neglect*, 153, 106811. DOI:10.1016/j.chiabu.2024.106811

Evidence overwhelmingly suggests that adverse childhood experiences (ACEs) is a risk factor for poor mental health outcomes. However, the specific mechanisms via which ACEs confer an increased risk of psychopathology are less well understood. The study modelled the effect of empathy & perceived social support (PSS) on mental health outcomes in a mixed clinical & non-clinical population, within the context of exposure to ACEs. A total of 575 participants aged 18 to 65 completed self-report measures assessing early adversity, PSS, empathy, and mental health outcomes. Multiple mediation analyses were used to investigate whether empathy and PSS mediated the relationship between self-reported ACEs & mental health outcomes, and whether affective & cognitive empathy affected differentially the link between emotional neglect & psychological distress. Results revealed a statistically significant indirect effect of ACEs on adult mental health through affective empathy and PSS. Emotional neglect was the only type of adversity significantly correlated with both dimensions of empathy. The indirect effect of emotional neglect on mental health outcomes via cognitive & affective empathy was also statistically significant. Study results highlight the role of affective empathy & PSS as transdiagnostic mechanisms influencing the pathway between early adversity & adult

mental health, and the importance of these when designing interventions aiming to promote well-being among those who have experienced childhood adversity.

Sattler, K., Yoon, S., & Lutolli, A. (2024). Trajectories of resilience among young children involved with child protective services. *Development and Psychopathology*, 36(1), 467–477. DOI:10.1017/S095457942200133X

Although child maltreatment is associated with short & long term maladaptive outcomes, some children are still able to display resilience. Currently, there is a limited understanding of how children's resilience changes over time after experiencing maltreatment, especially for young children. Therefore, the current study used a longitudinal, multidimensional approach to examine trajectories of resilience among very young children involved in child protective services and determine whether placement setting and caregiving behaviors are associated with resilience trajectories. This study used data from National Survey of Child and Adolescent Well-Being I and conducted repeated measures latent class analysis, focusing on children under 2 years old at baseline (n = 1,699). Results suggested that there were 3 trajectories of resilience: increasing resilience, decreasing resilience, and stable, low resilience. Caregiver cognitive stimulation was related to increasing trajectories of resilience compared to both decreasing & stable, low resilience. These findings illustrate the importance of caregiving behaviors for promoting resilience among a particularly vulnerable population.

Turgeon, J., Racine, N., McDonald, S., Tough, S., & Madigan, S. (2024). Maternal adverse childhood experiences, child resilience factors, and child mental health problems: A multi-wave study. *Child Abuse & Neglect*, 154, 106927. DOI:10.1016/j.chiabu.2024.106927

Research suggests that maternal ACEs have intergenerational consequences for offspring mental health. However, very few studies have investigated moderators of this

association. The objective of this longitudinal study was to examine whether child resilience factors moderated the association between maternal ACEs prior to age 18, and child-reported symptoms of anxiety, depression, hyperactivity, & inattention. The current study used data from 910 mother-child dyads. Participants were recruited in pregnancy from 2008 to 2010 as part of a longitudinal cohort study. Mothers had previously completed an ACEs questionnaire and reported on their child's resilience factors at child age 8-years. Children completed questionnaires about their mental health problems (symptoms of anxiety, depression, hyperactivity, and attention problems) at ages 10 and 10.5 years. Four moderation models were performed in total. Results revealed that maternal ACEs predicted child-reported symptoms of anxiety and depression. However, both these associations were moderated by higher levels of perceived child resilience factors. Specifically, there was no association between maternal ACEs and child mental health problems in the context of moderate and high levels of child resilience factors. Children who have the ability to solicit support from internal and external sources (e.g., being creative, setting realistic goals, making friends easily) may be buffered against the consequences of maternal ACEs on anxiety and depression. Thus, the effects of maternal ACEs on child mental health problems are not deterministic.

Bucci, S., Varese, F., Quayle, E., Cartwright, K., Machin, M., Whelan, P., Chitsabesan, P., Richards, C., Green, V., Norrie, J., & Schwannauer, M. (2023). [A digital intervention to improve mental health and interpersonal resilience in young people who have experienced technology-assisted sexual abuse: Protocol for a nonrandomized feasibility clinical trial and nested qualitative study](#). *JMIR Research Protocols*, 12, e40539. DOI:10.2196/40539

No evidence-based support has been offered to young people (YP) who have experienced technology-assisted sexual abuse (TASA). Interventions aimed at improving mentalization (the ability to understand the mental states of oneself and others) are increasingly being applied to treat YP with various clinical issues. Digital technology use

among YP is now common. A digital intervention aimed at improving mentalization in YP who have experienced TASA may reduce the risk of revictimization and future harm and make YP more resilient and able to manage distress that might result from TASA experiences. In this paper, we describe a protocol for determining the feasibility of the i-Minds trial and the acceptability, safety, and usability of the digital intervention (the i-Minds app) and explore how to best integrate i-Minds into existing routine care pathways. This is a mixed methods nonrandomized study aimed to determine the feasibility, acceptability, safety, and usability of the intervention. Participants aged between 12 and 18 years who report distress associated with TASA exposure will be recruited from the United Kingdom from the National Health Service (NHS) Trust Child and Adolescent Mental Health Services, sexual assault referral centers, and a web-based e-therapy provider. All participants will receive the i-Minds app for 6 weeks. Coproduced with YP and a range of stakeholders, the i-Minds app focuses on 4 main topics: mentalization, TASA and its impact, emotional and mental health, and trauma. A daily prompt will encourage YP to use the app, which is designed to be used in a stand-alone manner alongside routine care. We will follow participants up after the intervention and conduct interviews with stakeholders to explore the acceptability of the app and trial procedures and identify areas for improvement. Informed by the normalization process theory, we will examine barriers and enablers relevant to the future integration of the intervention into existing care pathways, including traditional clinic-based NHS and NHS e-therapy providers. This study was approved by the Research Ethics Board of Scotland. We expect data to be collected from up to 60 YP. We expect to conduct approximately 20 qualitative interviews with participants and 20 health care professionals who referred YP to the study. The results of this study have been submitted for publication. This study will provide preliminary evidence on the feasibility of recruiting YP to a trial of this nature and on the acceptability, safety, and usability of the i-Minds app, including how to best integrate it into existing routine care. The findings will inform the decision to proceed with a powered efficacy trial.

Lassri, D., Bregman-Hai, N., Soffer-Dudek, N., & Shahar, G. (2023). [The interplay between childhood sexual abuse, self-concept clarity, and dissociation: A resilience based perspective](#). *Journal of Interpersonal Violence*, 38(3-4), 2313-2336.
DOI:10.1177/08862605221101182

Despite a robust consensus regarding the potentially negative implications of childhood sexual abuse (CSA), research investigating risk & protective factors, particularly among well-functioning young adults, is scant. Dissociation is one of the major maladaptive outcomes of CSA but it only explains about 10% of the dissociation variance. Possibly, this modest effect size is due to protective factors moderating the relation between CSA and dissociative symptoms. One factor may be the extent to which one has succeeded in developing a clear & coherent sense of who they are. We aimed to explore whether self-concept clarity (SCC) moderates the relationship between CSA & dissociation (Model 1), and an alternative hypothesis, whereby CSA moderates the relationship between SCC and dissociation (Model 2). This was tested among young women, drawn from an earlier study that intentionally oversampled CSA survivors. We included data from survivors of CSA by a known perpetrator and women with no sexual trauma. Findings were consistent in both Model 1 & 2, but only when depersonalization-derealization, namely detachment, was considered. Simple effects analyses revealed that CSA was related to depersonalization-derealization only under low SCC levels (Model 1), and SCC was negatively related only in the CSA group (Model 2). Findings suggest that SCC is a protective factor, buffering the association between CSA and detachment symptoms.

Latham, R. M., Newbury, J. B., & Fisher, H. L. (2023). [A systematic review of resilience factors for psychosocial outcomes during the transition to adulthood following childhood victimisation](#). *Trauma, Violence, & Abuse*, 24(2), 946-965.
DOI:10.1177/15248380211048452

Exposure to childhood victimisation (i.e. abuse, neglect, domestic violence or bullying) can detrimentally impact later psychosocial adjustment. However, this is not the case for

all victimised children; some do well despite their experiences and are considered to be resilient. Understanding the factors associated with such resilience is important to inform interventions to support better psychosocial outcomes among victimised children. This review provides an overview of the extant research examining resilience factors for psychosocial outcomes during the transition to adulthood following exposure to childhood victimisation. Studies were identified through a systematic literature search of Embase, PsychINFO & Ovid MEDLINE databases. The 26 included studies spanned a range of psychosocial outcomes between ages 18–25, including education & work, housing & independent living, criminal behaviour, victimisation, and social & psychological adjustment. For each outcome, a variety of putative resilience factors had been investigated including those related to the individual, their family and the wider community within which they lived. However, because few studies had comparable resilience factors & psychosocial outcomes, it is difficult to draw conclusions about which factors are consistently associated with resilience to a particular psychosocial outcome. This review also revealed that the included studies were of variable methodological quality. Many were limited by cross-sectional designs with retrospective self-reports of childhood victimisation, and convenience or unrepresentative samples. In this review, we also highlight gaps in knowledge about the co-occurring impact of multiple resilience factors in combination and the need for studies conducted in non-Western, low- and middle-income countries.

Meraj, N., Arbeau, K., Fadiya, B., Ketelaars, T., St. Pierre, J., Swart, G. T., & Zayed, R. (2023). Introducing the Adverse Life Events Inventory for Children (ALEIC): An examination of adverse experiences and related impacts in a large clinical sample of children and youth. *Traumatology*, 29(2), 137–148. DOI:10.1037/trm0000385

Adverse childhood experiences (ACEs) investigators report a dose–response relation between the number of negative experiences in childhood and physical & emotional well-being in adulthood. Measures of ACEs have been developed for clinical use; however,

some instruments focus only on specific types of negative events and/or conflate trauma symptoms. We present the development of the Adverse Life Events Inventory for Children (ALEIC), a measure of childhood adversity exposure that enhances the comprehensive interRAI Child and Youth Mental Health intake instrument already in use at children's mental health agencies. ALEIC development was aided by an external Delphi panel of trauma experts. These efforts resulted in 28 items across 16 categories, covering a wide range of negative life events. Archival data analyzed from 13,645 children & youth who had initial Child and Youth Mental Health assessments found a positive relation between ALEIC total scores of adversity history and trauma mental health symptoms. As a benchmark for researchers & clinicians, we include prevalence rates reported by sex & age for each type of adversity in this large and diverse clinical sample. Remarkably, 40% of children and 51% of youth referred to mental health clinics experienced 4 or more (of 16) different types of adversity. Given these base rates, routine, structured screening of adversity exposure at clinic intake is recommended to mitigate the reoccurrence of future stressors and to minimize the long-term effects of adverse life events.

Pond, R., Gillmore, C., & Blanchard, N. (2023). Lived experiences of resilience for women who have experienced childhood sexual abuse: A systematic review of qualitative studies. *Child Abuse & Neglect*, 140, 106152.
DOI:10.1016/j.chiabu.2023.106152

Child sexual abuse (CSA) can have detrimental, long-lasting consequences but effects are diverse, and resilience, or attainment of better-than-expected outcomes, is possible. This systematic review synthesises qualitative research findings about lived experiences of resilience processes used by women who had been subjected to CSA. A comprehensive search was undertaken of major and minor article databases (e.g., PsychInfo, Medline, CINAHL, Web of Science, Scopus) and Google Scholar, with hand searching of reference lists and forward searching of retrieved articles. Eight studies met the inclusion criteria (peer-reviewed qualitative or mixed methods studies written in

English about experiences of resilience by women who had experienced CSA). Data extraction, quality appraisal, and thematic analysis were performed. Results: Thematic analysis generated several process-related resilience themes: distancing oneself from sexual abuse; building healthy interpersonal, community, and cultural connections; drawing on spiritual belief systems; reframing the sexual abuse; placing blame with the perpetrator; reclaiming self-worth; taking control of one's life; and pursuing meaningful goals. For some, it involved forgiveness of self and others, reclaiming one's sexuality, and/or fighting against diverse forms of oppression. There was much evidence that resilience was a dynamic, personal and social-ecological phenomenon. Conclusion: Counsellors and other professionals can use these findings to help women affected by CSA explore, develop, and strengthen factors that support resilience. Future research could explore the experiences of resilience for women with different cultural backgrounds, socioeconomic status, religious and/or spiritual affiliations.

Hébert, M., Amédée, L. M., Théorêt, V., & Petit, M. P. (2022). Diversity of adaptation profiles in youth victims of child sexual abuse. *Psychological Trauma: Theory, Research, Practice and Policy*, 14(S1), S41–S49. DOI:10.1037/tra0001090

Objective: Child sexual abuse (CSA) is associated with long-term negative consequences in adolescents, but some survivors display resilience. The purpose of this study was to delineate profiles of adaptation in adolescent victims of CSA and to examine their associations with individual and environmental-systemic protective factors. As part of a population-based survey, 8,230 high school students were questioned about CSA and completed measures assessing a host of protective factors and indicators of positive adaptation across 5 domains: self-perception, academic success, mental health, health risk behaviors and romantic relationships. Using a latent class analysis, a best fitting model of 4 classes was identified. This model included a reference group of nonsexually abused teenagers and 3 classes characterizing survivors of CSA: Resilient profile (33% of

youth), Externalized profile (34% of youth) and Internalized profile (33% of youth). Sexually abused youth assigned to the Resilient profile were similar to nonsexually abused youth in terms of self-esteem, academic performance, absence of clinical levels of psychological distress and dating violence. Despite experiencing CSA of comparable severity, youth in the Resilient profile reported more optimism and were less likely to rely on avoidant or emotional strategies to cope with difficulties and more likely to report high maternal and paternal support. Findings highlight the utility of a person-oriented approach to enhance our understanding of the diversity of adaptation profiles in youth victims of CSA. Results also underscore the importance of tailoring intervention efforts to efficiently tackle the diverse needs of teen victims of CSA.

Kumar, S. A., Brockdorf, A. N., Jaffe, A. E., Church, H. R., Messman, T. L., & DiLillo, D. (2022). [Mindful awareness promotes resilience: Buffered links among childhood sexual abuse severity, goal-directed emotion dysregulation, and psychopathology.](#) *Mindfulness*, 13(4), 993–1006. DOI:10.1007/s12671-022-01854-2

Objectives: Childhood sexual abuse is linked to long-term consequences, including depression and anxiety in adulthood. Although considerable progress has been made to understand mechanisms that may account for this relation, such as emotion dysregulation, less attention has been given to protective factors that may mitigate it. One such protective factor might be mindful awareness. Those who act with awareness in daily living tend to engage in healthy emotion regulation skills when faced with stressors and experience less depression and anxiety. In the current study, we aimed to replicate the positive associations among childhood sexual abuse severity, emotion dysregulation, and psychopathology across time, and also identify a personal strength—in this case, mindful awareness—that might mitigate these effects. Methods: Participants were 491 women recruited from the community who completed self-report assessments at three time points over a 32-month period. Results: A series of moderated mediation models revealed childhood sexual abuse severity predicted later reports of depression

and anxiety symptoms through greater emotion dysregulation in the form of difficulties engaging in goal-directed behaviors. As expected, mindful awareness weakened the relation between goal-directed emotion dysregulation and symptoms of depression and anxiety, such that greater levels of mindful awareness fully buffered these effects. Conclusions: Through a better understanding of natural resiliency processes among survivors, we can ultimately encourage continued examination of what might be effective additions to existing treatments for the mental health consequences of trauma and adversity.

Nuñez, M., Beal, S. J., & Jacquez, F. (2022). [Resilience factors in youth transitioning out of foster care: A systematic review](#). *Psychological Trauma: Theory, Research, Practice and Policy*, 14(S1), S72–S81. DOI:10.1037/tra0001096

Objective: Youth who emancipate from the foster care system often experience poor outcomes during their transition into independent living (e.g., criminal justice involvement and homelessness). Yet, some youth are resilient and achieve positive outcomes. The purpose of this study is to review the resilience factors identified in the literature for youth transitioning out of care. Method: A systematic review of the literature was conducted using PsycINFO, PubMed, and EMBASE databases. A total of 12 studies met our inclusion criteria. Results: A total of 38 different resilience factors were identified across the studies reviewed. From these resilience factors, 18 were statistically significant, including four assets and 14 resources. Conclusion: Understanding the assets and resources that promote resilience can aid clinicians in assessing and capitalizing on youths' strengths and can help researchers develop effective interventions and target gaps in the literature. Further, it can assist policymakers in generating legislation to improve outcomes for foster youth transitioning out of care. To assist in this process, we systematically reviewed studies to provide a summary of resilience factors currently identified in the literature.

Satapathy, S., Dang, S., Sagar, R., & Dwivedi, S. N. (2022). Resilience in children and adolescents survived psychologically traumatic life events: A critical review of application of resilience assessment tools for clinical referral and intervention. *Trauma, Violence, & Abuse, 23*(1), 288–300. DOI:10.1177/1524838020939126

Psychological traumatic life events (TLEs) and resilience, both are multidimensional, complex, and share salient features. Both are products of individual, familial, and environmental (socio-cultural-political contextual) variables, which is very crucial in children and adolescents. This systematic review used Boolean search strategies in electronic databases, namely, PubMed, PsycNET, JStor, and Google scholar. All researches not studying resilience per se but similar or related constructs such as life strengths, hardiness, protective/risk factors, social support, self-efficacy, social-emotional adjustment, and so on were excluded. A total of 12 resilience tools meant for children between 5 and 18 years were reviewed. The scale characteristics were analyzed in terms of targeted age-group of sample; purpose (i.e. screening and profiling for intervention); number of items; purpose/type of scale; year, country, and domain wise distribution; response format, standardization sample profile; psychometric properties; and availability of manual with norms of cutoff score. Although no scale was originally developed for children and adolescent population with history of TLEs particularly various forms of abuse and trauma, Child and Youth Resilience Measure and Connor-Davidson Resilience Scale had small samples of children from welfare homes. Neither did any scale tested the divergent validity against absence of any psychopathology or global functioning or poor quality of life Nor did Majority of scales provide a cutoff value for institutionalized children and adolescents with history of TLEs; therefore, using an existing scale for this purpose should be carefully examined. Trauma-focused multidimensionality in resilience needs to be explored more rigorously through mixed methods.

Yang, Y., Ma, X., Kelifa, M. O., Li, X., Chen, Z., & Wang, P. (2022). The relationship between childhood abuse and depression among adolescents: The mediating role of school connectedness and psychological resilience. *Child Abuse & Neglect*, 131, 105760. DOI:10.1016/j.chiabu.2022.105760

Background: Adolescent depression is a major public health concern worldwide, and depressive symptoms among adolescents in Chinese secondary school increase with grade. Although numerous studies noted that childhood abuse was associated with the development of depression in adolescents, the underlying mediators involved in this pathway remain unclear. Therefore, our study aims to explore whether psychological resilience and school connectedness mediate the association between childhood abuse and depression among Chinese adolescents. Methods: A cross-sectional survey was carried out among 1607 adolescents at four high schools located in Henan Province, Central China. Structural equation modeling was performed to determine whether psychological resilience and school connectedness mediate the influence of childhood abuse on depression. Results: Findings revealed that childhood abuse was directly ($\beta = 0.410, p < 0.001$) and indirectly ($\beta = 0.141, p < 0.001$) related to depression. Specifically, psychological resilience ($\beta = 0.018, p = 0.009$), school connectedness ($\beta = 0.087, p < 0.001$), and school connectedness together with psychological resilience ($\beta = 0.036, p < 0.001$) partially mediated the relationship between childhood abuse and depression. Conclusions: The detrimental impact of childhood abuse on adolescents' depression may be reduced by fostering psychological resilience and school connectedness.

Lo, J. (2021). [Fostering resilience in survivors of child sexual abuse to decrease vulnerability to sexual revictimization](#). *Crisis, Stress, and Human Resilience: An International Journal*, 3(3), 61–69.

Child Sexual Abuse (CSA) is an enormous public health issue worldwide, affecting health outcomes in millions, and the United States is not immune. One significant sequela of CSA

is increased vulnerability for sexual revictimization. Each CSA survivor has varying factors that affect their own risks for revictimization, and significant attempts to lower such risks must take into account their complexity. This author seeks to foster resilience in survivors of CSA, utilizing strengths and positive factors which are already part of the survivor as well as those who surround them, and by enhancing or teaching factors of resilience, to decrease sexual revictimization.

Yoon, S., Howell, K., Dillard, R., McCarthy, K. S., Napier, T. R., & Pei, F. (2021). Resilience following child maltreatment: Definitional considerations and developmental variations. *Trauma, Violence, & Abuse*, 22(3), 541–559.
DOI:10.1177/1524838019869094

Resilience following childhood maltreatment has received substantial empirical attention, with the number of studies on this construct growing exponentially in the past decade. While there is ample interest, inconsistencies remain about how to conceptualize and assess resilience. Further, there is a lack of consensus on how developmental stage influences resilience and how protective factors affect its expression. The current systematic review uses a developmental lens to synthesize findings on resilience following child maltreatment. Specifically, this article consolidates the body of empirical literature in a developmentally oriented review, with the intention of inclusively assessing three key areas—the conceptualization of resilience, assessment of resilience, and factors associated with resilience in maltreatment research. A total of 67 peer-reviewed, quantitative empirical articles that examined child maltreatment and resilience were included in this review. Results indicate that some inconsistencies in the literature may be addressed by utilizing a developmental lens and considering the individual's life stage when selecting a definition of resilience and associated measurement tool. The findings also support developmental variations in factors associated with resilience, with different individual, relational, and community protective factors emerging based on life stage. Implications for practice, policy, and research are incorporated throughout this review.

Gower, T., Rancher, C., Campbell, J., Mahoney, A., Jackson, M., McDonald, R., & Jouriles, E. N. (2020). Caregiver and divine support: Associations with resilience among adolescents following disclosure of sexual abuse. *Child Abuse & Neglect, 109*, 104681. DOI:10.1016/j.chiabu.2020.104681

This study examines relations of caregiver support and divine support with resilience following adolescent sexual abuse. Participants were 548 adolescents aged 11–17 (Mage = 13.78; 91% female) and their non-offending caregivers (Mage = 39.68; 79% mothers). Families were recruited from a children's advocacy center located in the southern United States following a disclosure of sexual abuse. Adolescents completed measures of caregiver and divine support and their social and academic functioning; caregivers and adolescents reported on adolescent psychological functioning. Resilience was operationalized as the absence of clinical levels of psychological symptoms together with the presence of adaptive levels of social and academic functioning. Both caregiver support ($r = .19, p < .001$) and divine support ($r = .14, p = .001$) were positively associated with resilience. Considered together in regression analyses, caregiver support (OR = 1.02, $p = .001$) and divine support (OR = 1.06, $p = .04$) were both associated with resilience. Clinicians and researchers should consider the potential contribution of caregiver and divine support to adolescent resilience following sexual abuse.

Hamby, S., Taylor, E., Mitchell, K., Jones, L., & Newlin, C. (2020). [Is it better to seek or to receive? A dual-factor model of social support](#). *International Journal of Child and Adolescent Resilience, 7*(1), 5–17. DOI:10.7202/1072584ar

This study adopts a dual-factor approach to examine the association of seeking and receiving social support with 6 indicators of current functioning and 14 psychosocial strengths. A survey completed by 440 youth ages 10 to 21 ($M = 16.38, SD = 3.04$) assessed strengths, functioning, and victimization. Youth were classified into four groups: Interconnected (high on social support seeking and receiving; 33% of sample), Rebuffed (high on social support seeking, low on social support receiving; 12%), Tended (low on

social support seeking, high on social support receiving; 16%), and Isolated (low on social support seeking and receiving; 39%). Controlling for age, gender, and victimization, the social support group was associated with each meaning making, regulatory, and interpersonal strength, and every indicator of current functioning except trauma symptoms. The Isolated group scored lowest on all measures and the Interconnected group scored highest on 19 of 20 measures. The mixed profile groups fell between these extremes. Notably, the Rebuffed group reported higher levels of some strengths and non-theistic spiritual well-being than the Tended group. The Tended group was never significantly higher than the Rebuffed group. Individual skills and attitudes regarding helpseeking may be more impactful than social support provided by others. Rebuffed youth may be steeling themselves in other strengths when the social environment is not supportive.

Pei, F., McCarthy, K. S., Wang, X., Beaujolais, B., & Yoon, S. (2020). Critical components of interventions to promote resilience among children with maltreatment experiences: A qualitative study of practitioners' perspectives. *Children and Youth Services Review, 110*, 104810. DOI:10.1016/j.chidyouth.2020.104810

Although researchers and practitioners keep working on identifying efficacious interventions for children with maltreatment experiences, few studies explore the perspective of the clinicians for their perspective of the critical components of interventions to promote resilience among maltreated children. To address this research gap, 27 interviews were conducted with clinicians to examine the critical components that should be incorporated into interventions. Four primary themes were captured from practitioners' perspective: (1) therapist characteristics; (2) caregiver characteristics; (3) community influences; and, (4) post-treatment support. Findings of this study benefit the improvement of future clinical interventions to promote the resilience of maltreated children.

Wamser-Nanney, R., & Campbell, C. L. (2020). Children's coping following sexual abuse: The roles of abuse characteristics, abuse stress, and maternal support. *Journal of Child & Family Studies*, 29, 514–525. DOI:10.1007/s10826-019-01540-3

Although the use of coping strategies is recognized as playing a vital role in predicting the development of trauma-related symptoms, little research has identified what factors may contribute to the utilization, and perceived efficacy of, coping strategies following childhood sexual abuse (CSA). The current study investigated whether CSA characteristics, abuse stress, and maternal support were related to children's utilization and perceived efficacy of active/social, internalized, angry, & avoidant coping strategies among 141 sexually abused children aged 7–12. Children used nearly 7 types of coping strategies. Longer CSA duration was associated with utilization of internalized & avoidant coping, and maternal blame/doubt predicted the use of avoidant coping. Girls utilized angry coping more frequently than boys. CSA characteristics and abuse stress were largely unrelated to children's perceived coping efficacy. Minority children reported higher levels of perceived efficacy of internalized strategies than white children. CSA duration may be important in understanding children's utilization of internalized & avoidant coping strategies and maternal blame/doubt may be related to children's use of avoidant coping. However, CSA characteristics and abuse stress may ultimately have little bearing on sexually abused children's use and perceived efficacy of coping strategies.

Yoon, S., Dillard, R., Pei, F., McCarthy, K. S., Beaujolais, B., Wang, X., MaGuire-Jack, K., Wolf, K., & Cochey, S. (2020). Defining resilience in maltreated children from the practitioners' perspectives: A qualitative study. *Child Abuse & Neglect*, 106, 104516. DOI:10.1016/j.chiabu.2020.104516

Increasingly, studies have focused on understanding positive outcomes in children who have been maltreated and the factors that contribute to resilience. However, there is no universally accepted definition of resilience, thus hindering the ability to make

comparisons across studies and to use such information to inform interventions to foster resilience. The current study sought to address this gap by examining definitions of resilience in practitioners who work directly with maltreated children. 27 participants were recruited through two agencies that serve victims of child maltreatment in an urban Midwestern city. Through a series of 27 qualitative interviews, the current study examined the following research question: "How is resilience defined and understood by practitioners working with children who have experienced child maltreatment?" Thematic coding and analysis were used to analyze the data. Findings suggest five unique themes described by practitioners as their definition of resilience: (a) surviving; (b) thriving; (c) perseverance; (d) reconciling and integrating traumatic experiences into healthy identity development; and (e) advocating for self.

Moisan, C. Hébert, M., Fernet, M., Blais, M., & Amédée, L. M. (2019). [Resilience portfolios and poly-strengths: Identifying strengths associated with wellbeing after adversity.](#) *International Journal of Child and Adolescent Resilience*, 6(1), 19–35.
DOI:10.7202/1069073ar

This study aimed to describe the prevalence of traumas and strengths in a representative sample of Quebec youth and to test whether poly-strengths were associated with low psychological distress, after controlling for poly-traumas. Using data from the Quebec Youths' Romantic Relationships survey (QYRRS), hierarchical logistic regressions were conducted to examine the relationship between poly-strengths and low levels of psychological distress, and to identify which strengths were associated with outcomes, after accounting for demographic variables and individuals' experiences of traumas. More than a third of the sample experienced 4 traumas or more (37.0%). The average number of experienced traumas was 3.04 out of 10 measured traumas. More than half of the sample had at least 5 strengths, the average number of strengths being 3.95 (out of 8). Two third (67.6%) of the sample did not suffer from psychological distress. Among poly-victims, half of the participants (49.6%) showed clinical symptoms of distress. Poly-

strengths were uniquely associated with low of clinical distress. After accounting for demographics and poly-traumas, poly-strengths explained 24.2% of the variance of low levels of psychological distress. Self-esteem, optimism, parental support and attachment, number of sources of support, social support (seeking secure base), and capacity to adapt (resiliency) were uniquely associated with low levels of distress. The combination of strengths decreases the likelihood of experiencing clinical levels of psychological distress, which can contribute to healthy functioning in context of adversities. Findings highlight the importance of promoting multiple and diverse strengths among youth.

Hamby, S., Taylor, E., Jones, L., Mitchell, K. J., Turner, H. A., & Newlin, C. (2018). From poly-victimization to poly-strengths: Understanding the web of violence can transform research on youth violence and illuminate the path to prevention and resilience. *Journal of Interpersonal Violence*, 33(5), 719-739.
DOI:10.1177/0886260517744847

For many years, an overly “siloe” approach has hampered efforts to understand violence and minimize the societal burden of violence and victimization. This article discusses the limitations of an overly specialized approach to youth violence research, which has focused too much on violence in particular contexts, such as the family or the school. Instead, a child-centered approach is needed that comprehensively assesses all exposures to violence. This concept of the total cumulative burden of violence is known as poly-victimization. The poly-victimization framework reveals that many youth are entangled in a web of violence, experiencing victimization in multiple settings by multiple perpetrators. This more accurate view of children’s exposure to violence has many advantages for advancing our scientific understanding of violence. Perhaps somewhat surprisingly, this more comprehensive view also points to new insights for resilience and prevention. This includes recognizing a parallel concept, “poly-strengths,” which captures the number of resources and assets children and their families can use to help insulate

youth from violence (prevention) or assist in coping and promoting well-being after victimization (intervention). Reconceptualizing how resilience is defined and understood among youth populations can help alleviate the true societal burden of youth victimization.

Ressel, M., Lyons, J., & Romano, E. (2018). Abuse characteristics, multiple victimization and resilience among young adult males with histories of childhood sexual abuse. *Child Abuse Review*, 27(3), 239–253. DOI:10.1002/car.2508

Although researchers now concede that one in six men experiences childhood sexual abuse (CSA), this population remains understudied in both the empirical and clinical literature. Little is known about the characteristics of males' CSA experiences (e.g. duration, child-perpetrator relationship) and about resilient functioning. This study described young adult males' abuse experiences during childhood and their current adaptive functioning. Participants were recruited from across North America through websites geared specifically for males with CSA histories. The sample included 46 males aged 17–25 years who anonymously completed an online questionnaire. Findings indicated that males tended to experience severe CSA, including early age of onset, invasive sexual acts and the use of perpetrator force. Males also reported the co-occurrence of other forms of childhood maltreatment and adversity, including parental conflict, neglect, physical abuse and emotional maltreatment. Finally, males reported lower rates of resilience on standardised measures, compared with adult community samples in North America. However, scores were consistent with clinical age-based norms for adolescents with trauma histories. This study contributes to the growing awareness of male CSA, multiple victimisation and its diverse outcomes. Implications of these findings for future research and clinical practice are considered.

Sattler, K. M., & Font, S. A. (2018). [Resilience in young children involved with child protective services](#). *Child Abuse & Neglect*, 75, 104–114.
DOI:10.1016/j.chiabu.2017.05.004

Child maltreatment increases the risk of poor developmental outcomes. However, some children display resilience, meaning they are high-functioning despite their adverse experiences. To date, few research studies have examined protective factors among very young maltreated children. Yet, levels of resilience, and the protective factors that promote resilience among maltreated children, are likely to differ by developmental stage. Drawing on ecological systems theory and life course theory, we examined how protective factors at multiple ecological levels across early childhood were related to social and cognitive resilience among very young children involved with child protective services. The results demonstrated that the buffering effects of protective factors varied by social or cognitive resilience and the cumulative effects of protective factors were more consistently related to later resilience than protective factors at specific time points. In addition, the influence of specific protective factors on resilience slightly varied by initial in-home or out-of-home placement. These findings have important policy and research implications for promoting optimal development among children involved in child protective services.

Happer, K., Brown, E. J., & Sharma-Patel, K. (2017). Children's resilience and trauma specific cognitive behavioral therapy: Comparing resilience as an outcome, a trait, and a process. *Child Abuse & Neglect*, 73, 30–41.
DOI:10.1016/j.chiabu.2017.09.021

Resilience, which is associated with relatively positive outcomes following negative life experiences, is an important research target in the field of child maltreatment. The extant literature contains multiple conceptualizations of resilience, which hinders development in research and clinical utility. Three models emerge from the literature: resilience as an immediate outcome (i.e., behavioral or symptom response), resilience as a trait, and

resilience as a dynamic process. The current study compared these models in youth undergoing trauma-specific cognitive behavioral therapy. Results provide the most support for resilience as a process, in which increase in resilience preceded associated decrease in posttraumatic stress and depressive symptoms. There was partial support for resilience conceptualized as an outcome, and minimal support for resilience as a trait. Results of the models are compared and discussed in the context of existing literature and in light of potential clinical implications for maltreated youth seeking treatment.

Kassis, W., Artz, S., Maurovic, I., & Simões, C. (2017). What doesn't kill them doesn't make them stronger: Questioning our current notions of resilience. *Child Abuse & Neglect*, 78, 71-84. DOI:10.1016/j.chiabu.2017.12.011

This study expands on earlier analyses of the data generated by a cross-sectional study involving a random sample of 5149 middle-school students with a mean age of 14.5 years from 4 EU countries (Austria, Germany, Slovenia, and Spain), in which every fourth respondent had been physically abused by his or her parents and almost every sixth respondent had witnessed physical spousal abuse. Contrary to expectations, some of these youths reported no engagement in peer violence and no symptoms of depression, which meant that they could be considered "resilient." Given their precarious conditions, we inquired into how these young people functioned on other protective and risk indicators when compared to non-violence exposed peers. Using Bonferroni post-hoc tests, we conducted an analysis of variance based comparison of levels of risk and protective factors on three groups of violence and depression-resilient youth (low, middle and high family violence experience) with those participants who reported no family violence or abuse, no depression and no use of violence. The violence and depression-resilient participants reported significantly higher levels of aggression supportive beliefs, alcohol consumption, drug use, verbal aggression towards and from teachers and use of indirect aggression, along with lower levels of social & personal protective characteristics

such as self- acceptance, emotional self-control, optimism about the future, and positive relations with parents & teachers, than students without family violence experiences. We therefore concluded that while some family violence exposed young people may not engage in violence or experience depression, this does not automatically imply an absence of other challenges and calls into question our current notions of resilience.

Newsom, K., & Myers-Bowman, K. (2017). "I am not a victim. I am a survivor": Resilience as a journey for female survivors of child sexual abuse. *Journal of Child Sexual Abuse*, 26(8), 927-947. DOI:10.1080/10538712.2017.1360425

This study was designed to expand our understanding of the positive aspects of coping & resilience in female survivors of child sexual abuse (CSA). Research questions focused on women's lived experiences of being survivors of CSA and how they have experienced resilience, developed healthy intimate relationships, and viewed themselves as sexual beings. Using a qualitative research lens of phenomenology, we captured the essence of survivors' experiences of resilience. Although each woman's experience was unique, similar patterns of processes & outcomes emerged as meaningful in their development of healthy sexuality and relationship functioning. Participants emphasized resilience as a process or journey—recovery from trauma, reconceptualization of self, and development of healthy sexuality included deliberate efforts occurring over time. Implications for future research and practice using a positive lens of resilience are provided.

Pérez-González, A., Guilera, G., Pereda, N., & Jarne, A. (2017). Protective factors promoting resilience in the relation between child sexual victimization and internalizing and externalizing symptoms. *Child Abuse & Neglect*, 72, 393-403. DOI:10.1016/j.chiabu.2017.09.006

Sexual victimization has been one of the most frequently studied forms of child victimization. Its effects are common and diverse; however, not all children and youth

exposed to sexual victimization eventually develop adjustment problems. A total of 1105 children and youth (590 male and 515 female) aged between 12 and 17 from northeastern Spain were assessed regarding their experiences of sexual victimization, symptoms of psychopathology, and protective factors. The results showed that all forms of sexual victimization were associated with higher levels of emotional and behavioral problems. However, the presence of a low Negative Cognition, high Social Skills and high Confidence seem to act buffering internalizing problems. Additionally, a significant interaction between Sexual Victimization and low Negative Cognition was observed ($p < 0.5$), so that, low Negative Cognition was related to a lower risk of being in the clinical range for internalizing problems. Likewise, high scores on Empathy/Tolerance, Connectedness to School, Connectedness to Family and low Negative Cognition acted as promotive factors in relation to externalizing symptoms, in this case without any interaction effect. The strong relationship found with emotional and behavioral problems highlights the importance of continuing the research on the protective factors underlying resilience in the relationship between sexual victimization and psychopathological symptoms. The findings also support the multi-dimensional and specific nature of resilience and identify some of the protective factors that should be regarded as key intervention targets in adolescents with a history of sexual victimization.

Day, T., & Kearney, C. A. (2016). Resilience variables and posttraumatic symptoms among maltreated youth. *Journal of Aggression, Maltreatment & Trauma*, 25(9), 991-1005. DOI:10.1080/10926771.2016.1236358

Child maltreatment is associated with an array of social, behavioral, neurobiological, and developmental problems that can last for many years. A key outcome of child maltreatment includes posttraumatic stress symptoms such as reexperiencing, avoidance, and hyperarousal. Considerable research has focused on risk factors for posttraumatic symptoms in maltreated youth, but less so on possible protective factors

such as resilience. This study examined several resilience variables (sense of mastery, sense of relatedness, emotional reactivity) in an ethnically diverse sample of maltreated adolescents with posttraumatic symptoms. Key aspects of resilience that predicted posttraumatic stress disorder (PTSD) symptoms included sense of mastery (for fewer symptoms) and emotional reactivity (for more symptoms). In particular, optimism, self-efficacy, and adaptability were inversely related to PTSD symptoms. In addition, emotional reactivity moderated a relationship between resilience resource scores and PTSD symptoms. The findings appeared to apply most to sexually maltreated youth. The results preliminarily demonstrate the importance of considering resilience-based variables during clinical processes for maltreated youth.

Dennison, M. J., Sheridan, M. A., Busso, D. S., Jenness, J. L., Peverill, M., Rosen, M. L., & McLaughlin, K. A. (2016). [Neurobehavioral markers of resilience to depression amongst adolescents exposed to child abuse](#). *Journal of Abnormal Psychology*, 125(8), 1201-1212. DOI:10.1037/abn0000215

Childhood maltreatment is strongly associated with depression, which is characterized by reduced reactivity to reward. Identifying factors that mitigate risk for depression in maltreated children is important for understanding etiological links between maltreatment and depression as well as improving early intervention and prevention. We examine whether high reward reactivity at behavioral and neurobiological levels is a marker of resilience to depressive symptomology in adolescence following childhood maltreatment. A sample of 59 adolescents (21 with a history of maltreatment; Mean Age 16.95 years, SD 1.44) completed an fMRI task involving passive viewing of emotional stimuli. BOLD signal changes to positive relative to neutral images were extracted in basal ganglia regions of interest. Participants also completed a behavioral reward-processing task outside the scanner. Depression symptoms were assessed at the time of the MRI and again 2 years later. Greater reward reactivity across behavioral and neurobiological measures moderated the association of maltreatment with baseline depression.

Specifically, faster reaction time (RT) to cues paired with monetary reward relative to those unpaired with reward and greater BOLD signal in the left pallidum was associated with lower depression symptoms in maltreated youth. Longitudinally, greater BOLD signal in the left putamen moderated change in depression scores over time, such that higher levels of reward response were associated with lower increases in depression over time among maltreated youths. Reactivity to monetary reward and positive social images, at both behavioral and neurobiological levels, is a potential marker of resilience to depression among adolescents exposed to maltreatment. These findings add to a growing body of work highlighting individual differences in reactivity to reward as a core neurodevelopmental mechanism in the etiology of depression.

Hamby, S., Banyard, V., & Grych, J. (2016). Strengths, narrative, and resilience: Restorying resilience research. *Psychology of Violence*, 6(1), 1-7. DOI:10.1037/vio0000027

Objective: To envision a path toward a more strengths-based approach to violence research, prevention, and intervention—a path that focuses on thriving and resilience. Key Points: Both the content and the process of research need to change if we are to transform our efforts to understand and overcome adversity. Greater focus on strengths and the achievement of well-being despite adversity is 1 important avenue; focusing on the narrative and the power of story is another important path. However, merely shifting the focus of traditional research and scholarly efforts is not enough. At another level of analysis, the field needs communication across the fragmentary subdisciplines of social science (“silo busting,” as we informally call it). We must also do more to encourage experimentation and innovation with regard to research question and design, community–practitioner–researcher partnership, and approaches to dissemination. Implications: Existing challenges in innovation and experimentation call for trying new approaches. Specific suggestions for adapting conference formats are provided. The commentaries in this special section offer feasible actions that could improve violence

research, including incorporating measures of well-being in addition to symptoms as outcome measures; involving a wider variety of stakeholders in research design and dissemination; taking advantage of new insights from positive psychology and narrative research; and incorporating aspects of community and culture into research, assessment, prevention and intervention.

Tlapek, S. M., Auslander, W., Edmond, T., Gerke, D., Voth Schrag, R., & Threlfall, J. (2016). The moderating role of resiliency on the negative effects of childhood abuse for adolescent girls involved in child welfare. *Children and Youth Services Review*, 73, 437–444. DOI:10.1016/j.childyouth.2016.11.026

Child abuse is associated with a number of negative psychosocial outcomes, including mental health problems, substance abuse, and interpersonal violence. Yet some children thrive or “adapt well” despite experiencing abuse; researchers are increasingly interested in the protective factors and mechanisms that may prevent negative outcomes. The purpose of this study was to explore the role of resiliency characteristics in the relationship between childhood abuse and the negative outcomes of depression, posttraumatic stress disorder (PTSD), substance abuse, and revictimization in adolescent girls involved in child welfare. Participants were 237 adolescent girls ages 12–19 years, and who were youths of color (75%) and white (25%). Data were collected through surveys assessing histories of physical, sexual, and emotional child abuse; symptoms of posttraumatic stress & depression; lifetime substance use; and experiences of physical, verbal, and relational revictimization in the last 3 months. All forms of childhood abuse were significantly associated with symptoms of depression and PTSD as well as revictimization and substance use. Higher levels of resiliency were associated with fewer depression and PTSD symptoms and experiencing less revictimization. Resiliency significantly moderated the relationship between sexual abuse & depression, PTSD, and revictimization. Resiliency also moderated the relationship between emotional abuse and

depression. No significant interactions emerged in the relationship between physical abuse and any of the mental health & behavioral problems.

Domhardt, M., Münzer, A., Fegert, J. M., & Goldbeck, L. (2015). Resilience in survivors of child sexual abuse: A systematic review of the literature. *Trauma, Violence, & Abuse, 16*(4), 476–493. DOI:10.1177/1524838014557288

This review article summarizes empirical research on resilience in survivors of child sexual abuse (CSA) and discusses protective factors that are associated with adaptive functioning in spite of sexual victimization. A literature search to identify studies published up to November 2013 was performed within the databases PsycINFO, MEDLINE/PubMed, Web of Science, & PSYINDEXplus. Additional relevant studies were retrieved using a snowball technique. A total of 37 articles met the inclusion criteria and were included in the final sample. In the studies included in this review, the percentage of CSA survivors who were found to have a normal level of functioning despite a history of sexual abuse ranged from 10% to 53%. The protective factors that had the best empirical support were found to be education, interpersonal & emotional competence, control beliefs, active coping, optimism, social attachment, external attribution of blame, and most importantly, support from the family & the wider social environment. Preventive and clinical interventions for survivors of CSA should utilize psychoeducation and cognitive strategies that are adapted to the developmental level of the victim and that seek to enhance social support from significant others. Future research should focus on longitudinal research designs considering resilience rather as a dynamic process with multiple dimensions in a social & developmental context.

Flett, G. L., Flett, A. L., & Wekerle, C. (2015). [A conceptual analysis of interpersonal resilience as a key resilience domain: Understanding the ability to overcome child sexual abuse and other adverse interpersonal contexts](#). *International Journal of Child and Adolescent Resilience*, 3(1), 4-33.

We conduct a conceptual analysis of interpersonal resilience as a domain-specific type of resilience, based on the premise that it is a multi-faceted construct. We consider interpersonal resilience within the context of child sexual abuse (CSA) as an under-attended and salient interpersonal stressor with profound implications involving the self & personal identity. Undoubtedly the most under-reported form of abuse, we examine the statistics known-to-date to highlight urgent areas for attention, pressing for resilience and developmentally focused empirical investigation. We describe interpersonal resilience as a developed orientation that is deeply rooted in self and identity issues. Interpersonal resilience incorporates processes that develop a sense of interpersonal efficacy, social self-esteem, mattering, and self-compassion that may buffer against negative social experiences, specifically the traumatic event of CSA embedded within adverse contexts. Interpersonal resilience is a distinct type of resilience, distinguishable from emotional resilience & dispositional traits. Empirical research on the nature of interpersonal resilience in challenging contexts is warranted. Intervention programs need to be expanded to include an explicit emphasis on practical resilience strategies, including promoting interpersonal resilience through skill-development, mentoring, and community-based opportunities.

Grych, J., Hamby, S., & Banyard, V. (2015). The resilience portfolio model: Understanding healthy adaptation in victims of violence. *Psychology of Violence*, 5(4), 343-354. DOI:10.1037/a0039671

Exposure to diverse forms of interpersonal violence is associated with a wide range of psychological problems in children and adults. However, many people who experience violence do not develop symptoms of psychopathology. Studies of resilience in victims of

violence have identified protective factors associated with healthier outcomes but have a number of limitations for understanding how individuals exposed to violence adapt and even thrive. The present article addresses these limitations by introducing a conceptual framework that integrates insights from theory and research on resilience, positive psychology, posttraumatic growth, and stress and coping. The Resilience Portfolio Model is a strengths-based framework designed to provide a holistic understanding of the protective factors and processes that promote resilience in children and adults exposed to violence. It proposes that the density and diversity of resources and assets available to individuals (their resilience portfolio) shapes their responses to violence and identifies 3 higher-order functional categories of strengths that are proposed to be particularly salient for resilience: regulatory, interpersonal, and meaning-making strengths. The Resilience Portfolio Model offers new directions for studying resilience in victims of violence and identifies a wider range of strengths and protective factors to address in prevention and intervention efforts.

Holmes, M. R., Yoon, S., Voith, L. A., Kobulsky, J. M., & Steigerwald, S. (2015). [Resilience in physically abused children: Protective factors for aggression](#). *Behavioral Sciences*, 5(2), 176–189. DOI:10.3390/bs5020176

Aggression continues to be a serious problem among children, especially those children who have experienced adverse life events such as maltreatment. However, there are many maltreated children who show resilient functioning. This study investigated potential protective factors (*i.e.*, child prosocial skills, child internalizing well-being, and caregiver well-being) that promoted positive adaptation and increased the likelihood of a child engaging in the healthy, normative range of aggressive behavior, despite experiencing physical maltreatment. Logistic regression analyses were conducted using two waves of data from the National Survey of Child and Adolescent Well-Being (NSCAW-I). Children who were physically maltreated were more likely to exhibit clinical levels of

aggressive behavior at Time 1 than children who were not physically maltreated. Children's internalizing well-being, children's prosocial behavior, and caregivers' well-being were associated with lower likelihood of clinical levels of aggressive behavior at Time 1. Children's internalizing well-being and children's prosocial behavior remained significantly associated with nonclinical aggression 18 months later. These findings highlight the role of protective factors in fostering positive and adaptive behaviors in maltreated children. Interventions focusing on preventing early aggression and reinforcing child prosocial skills, child internalizing well-being, and caregiver well-being may be promising in promoting healthy positive behavioral adjustment.

McGuigan, W. M., & Stephenson, S. J. (2015). A single-case study of resiliency after extreme incest in an Old Order Amish family. *Journal of Child Sexual Abuse*, 24(5), 526–537. DOI:10.1080/10538712.2015.1042182

This exploratory research brief presents a single case study of the resiliency of “Mary B.” She grew up in an Old Order Amish family where isolation, secrecy, and patriarchy masked repeated sexual assaults by her older brothers that began at age 7. By the age of 20, Mary alleged she had been raped on more than 200 separate occasions by members of her Amish family. After years of pleading with her mother and church officials to intervene, she sought therapy outside the Amish community. This led to three of her brothers being incarcerated. Her family disowned her and she was banned from the Amish community, leaving with an 8th grade education and little more than the clothes she was wearing. In less than 2 years, Mary had moved to a new town, completed her GED, obtained a car and driving license, maintained a small home, and worked as a certified nursing assistant. She consented to tape recorded interviews and completed several quantitative diagnostic measures. Scores on the diagnostic measures placed her within the normal range on self-esteem, competency, depression, stress, social support, and life skills. Analysis of interviews revealed Mary rebounded from her past by reframing her

experiences. Themes identified within the interviews supported 6 of the 7 types of resiliencies (insight, independence, initiative, relationships, humor, and morality) outlined in the therapeutic Challenge Model.

Hébert, M., Langevin, R., & Charest, F. (2014). [Factors associated with resilience in preschoolers reporting sexual abuse: A typological analysis](#). *International Journal of Child and Adolescent Resilience*, 2(1), 46–58.

The objectives of this study were to explore the diversity of profiles in sexually abused preschoolers and identify possible protective factors associated with individual differences in outcomes. A sample of 68 sexually abused children (ages 3½ – 6 ½ years old) and a comparison group of 78 children participated in the study. Parents evaluated children's level of internalizing and externalizing behavior problems with the Child Behavior Checklist (CBCL; Achenbach & Rescorla, 2000; 2001). They also reported on within-child protective factors by completing the Devereux Early Childhood Assessment (DECA; LeBuffe & Naglieri, 1999), their resilience capacity (CD-RISC 10; Campbell-Sills & Stein, 2007) and coping strategies (WOC; Folkman & Lazarus, 1988). A two-step cluster analysis was used to identify relevant subgroups of children. A three-cluster solution identified: a) High symptomatology subgroup whose members had clinically elevated scores on internalizing and externalizing behavior problems scales; b) moderate symptomatology group displaying significant externalizing behavior problems when compared to non-abused children, and c) resilient group of children displaying few behavior problems and benefiting from a host of protective factors. Results underscore the relevance of incorporating screenings for protective factors in addition to behavioral concerns in the assessments of sexually abused preschool-age children. Such an approach is likely to optimize the implementation of interventions for this vulnerable population.

Marriott, C., Hamilton-Giachritsis, C., & Harrop, C. (2014). Factors promoting resilience following childhood sexual abuse: A structured, narrative review of the literature. *Child Abuse Review*, 23(1), 17–34. DOI:10.1002/car.2258

The aim of this paper was to review research investigating resilient outcomes for people with a history of childhood sexual abuse (CSA) and implications for practice, as well as to consider issues for clearer definitions. Fifty English language peer-reviewed studies (1991–2010) met the inclusion criteria. The reviewed papers identified a number of factors that were repeatedly associated with individuals showing resilient outcomes to CSA. These included inner resources (e.g. coping skills, interpretation of experiences and self-esteem), family relationships, friendships, community resources (e.g. church or school), as well as some abuse-related factors (e.g. older age at onset). A large number of methodological concerns within these studies were also noted, including the way in which resilience, CSA and protective factors were defined. However, despite this, many papers identified similar factors that could be utilised to develop both effective prevention programmes and resilience interventions for the survivors of CSA.

Barnett, D., Heinze, H. J., & Arble, E. (2013). Risk, resilience, and the Rorschach: A longitudinal study of children who experienced sexual abuse. *Journal of Personality Assessment*, 95(6), 600–609. DOI:10.1080/00223891.2013.823437

Experiencing sexual abuse increases the risk that children will report or otherwise demonstrate problems with emotion, behavior, and health. This longitudinal study of 44 children who experienced sexual abuse examined whether information processing as assessed via the Rorschach Inkblot Test was associated with child-reported depression symptoms assessed via the Children's Depression Inventory (Kovacs, 1992) concurrently and an average of 15 months later. Children whose Rorschach protocols were relatively free of scores suggesting intense distress, complex processing, and sexual content were more likely to experience remission of depression symptoms at follow-up. Findings

provide incremental validity for certain Rorschach indexes to inform prognosis regarding depression symptoms and perhaps their treatment.

Cicchetti, D. (2013). [Annual research review: Resilient functioning in maltreated children Past, present, and future perspectives](#). *Journal of Child Psychology and Psychiatry*, 54(4), 402–422. DOI:10.1111/j.1469-7610.2012.02608.x

Through a process of probabilistic epigenesis, child maltreatment progressively contributes to compromised adaptation on a variety of developmental domains central to successful adjustment. These developmental failures pose significant risk for the emergence of psychopathology across the life course. In addition to the psychological consequences of maltreatment, a growing body of research has documented the deleterious effects of abuse and neglect on biological processes. Nonetheless, not all maltreated children develop maladaptively. Indeed, some percentage of maltreated children develops in a resilient fashion despite the significant adversity and stress they experience. Methods: The literature on the determinants of resilience in maltreated children is selectively reviewed and criteria for the inclusion of the studies are delineated. Results: The majority of the research on the contributors to resilient functioning has focused on a single level of analysis and on psychosocial processes. Multilevel investigations have begun to appear, resulting in several studies on the processes to resilient functioning that integrate biological/genetic and psychological domains. Conclusions: Much additional research on the determinants of resilient functioning must be completed before we possess adequate knowledge based on a multiple levels of analysis approach that is commensurate with the complexity inherent in this dynamic developmental process. Suggestions for future research on the development of resilient functioning in maltreated children are proffered and intervention implications are discussed.

Topitzes, J., Mersky, J. P., Dezen, K. A., & Reynolds, A. J. (2013). Adult resilience among maltreated children: A prospective investigation of main effect and mediating models. *Children and Youth Services Review*, 35(6), 937–949.
DOI:10.1016/j.chilyouth.2013.03.004

Studies examining resilience to child maltreatment reveal that maltreatment victims can achieve adaptive functioning in several areas of development; however, few of these individuals persistently demonstrate resilience across multiple domains. The majority of these investigations define adjustment with a limited number of outcomes measured proximal in time to the maltreatment experience. In contrast, this study measured adjustment across a diverse set of domains during early adulthood (ages 16–24), a number of years after the occurrence of childhood maltreatment (ages 0–11). Main effect & mediation analyses were conducted. Data were derived from the Chicago Longitudinal Study, an examination of 1539 minority individuals born in 1979 or 1980 in low-income Chicago-area neighborhoods. Study participants were followed prospectively from birth through age 24. Maltreatment data originated from official court & child protective service records. Parent report, self-report, and administrative sources informed covariate, mediator & outcome measures. Results from multivariate probit regression revealed that childhood maltreatment significantly & negatively predicted adult resilience. Exploratory and confirmatory mediation analyses showed the indicators that helped explain the long-term association between childhood maltreatment and young adult adjustment: school moves & out-of-home placement, reading ability, social skills, acting out behavior, juvenile delinquency, school commitment, and expectation to attend college.

Ungar, M. (2013). Resilience after maltreatment: The importance of social services as facilitators of positive adaptation. *Child Abuse & Neglect*, 37(2–3), 110–115.
DOI:10.1016/j.chiabu.2012.08.004

This practice note will show that resilience among children who have been maltreated is the result of multiple protective factors, including the quality of services provided to

children exposed to chronic adversity. This social ecological perspective of resilience suggests that it is a process resulting from interactions between individuals & their environments, dependent upon individual characteristics (personality & temperament), the social determinants of health affecting children and their families, formal interventions by multiple service providers (child welfare, special education, mental health, addictions, public health, & juvenile corrections), and the social policies that influence service provision to vulnerable populations. Clinicians & researchers concerned with the resilience of chronically abused & neglected children have tended to overlook the protective processes unique to children who have been abused that are different from the protective processes that promote positive development among children who have experienced no maltreatment. Most importantly, studies of resilience among maltreated children have rarely investigated the impact child welfare interventions have on the resilience of maltreated children, mistakenly attributing children's abilities to cope to be the result of individual factors rather than the responsiveness of service providers and governments to tailor interventions to children's needs. To enhance the likelihood of resilience among maltreated children, those who design and implement interventions need to address three aspects of resilience-related programming: make social supports & formal services more available & accessible; design flexible programs that can respond to the differential impact specific types of interventions have on children exposed to different forms of maltreatment; and design interventions to focus on subpopulations of children who experience maltreatment rather than diffuse population-wide initiatives.

Wekerle, C. (2013). Resilience in the context of child maltreatment: Connections to the practice of mandatory reporting. *Child Abuse & Neglect*, 37(2-3), 93-101.
DOI:10.1016/j.chiabu.2012.11.005

A human rights perspective places the care for children in the obligation sphere. The duty to protect from violence is an outcome of having a declaration confirming inalienable

human rights. Nationally, rights may be reflected in constitutions, charters, and criminal codes. Transnationally, the United Nation's (UN) Convention on the Rights of the Child (CRC) prioritizes a child's basic human rights, given their dependent status. Laws requiring professionals to report child maltreatment to authorities is one practical strategy to implement minimal child protection and service standards. Mandatory reporting laws officially affirm the wrong of maltreatment and the right of children and can be conceptualized as part of a resilience process, where the law sets the stage for child safety & well-being planning. Although widely enacted law, sizeable research gaps exist in terms of statistics on mandatory reporting compliance in key settings; obstacles & processes in mandatory reporting; the provision of evidence-based training to support the duty to report; and the training-reporting-child outcomes relationship, this latter area being virtually non-existent. The fact that mandatory reporting is not presently evidence-based cannot be separated from this lack of research activity in mandatory reporting. Reporting is an intervention that requires substantial inter-professional investment in research to guide best practices, with methodological expectations of any clinical intervention. Child abuse reporting is consistent with a clinician's other duties to report (suicidality, homicidality), practice-based skills (delivering "bad" news, giving assessment feedback), and the pervasive professional principle of best interests of the child. Resilience requires the presence of resources and, mandated reporting, is one such resource to maltreated children. Practice strategies identified in the literature are discussed.

Wekerle, C. (2013). [Practical strategies-maltreated adolescents: social and health service opportunities for resilience](#). *International Journal of Child and Adolescent Resilience*, 1(1), 88-96.

Well-being is well doing. The United Nations Rights on the Convention of the Child (CRC) and the United Nations Development Program Millennium Development Goals (MDGs)

target youth health, safety, participation and empowerment as key driver's of living in a social justice context. Health and social services are two streams of resources to at-risk youth and vulnerable caregivers to promote resilience – positive adaptation and development – in adverse contexts. Child maltreatment statistics highlight the critical role of the social safety net. A policy-service gap exists for youth involved in the Child Protective Services (CPS) system. For these youth, violence prevention and personal safety remains a key component of resilience. This *practical strategies* paper considers youth resilience issues the context of maltreatment. Innovation in adolescent protection is an opportunity for health and social service systems to support resilience, continuity of care and violence prevention. Strategies include transition services for the switch from child to adult services, prevention programming that support mediating the environment, targeting CPS expectant teens for established maltreatment prevention programming, and system changes that recognize the emerging adulthood developmental period.

Stumblingbear-Riddle, G., & Romans, J. S. (2012). Resilience among urban American Indian adolescents: Exploration into the role of culture, self-esteem, subjective well-being, and social support. *American Indian and Alaska Native Mental Health Research: The Journal of the National Center*, 19(2), 1-19.
DOI:10.1016/j.chiabu.2012.11.005

The effects of enculturation, self-esteem, subjective well-being, and social support on resilience among urban American Indian (AI) adolescents from a South Central region of the U.S. were explored. Of the 196 participants, 114 (58.2%) were female and 82 (41.8%) were male (ages 14-18 years). Thirty-three percent of the variance in resilience was accounted for by enculturation, self-esteem, and social support, while 34% of the variance in resilience was contributed by enculturation, subjective well-being, and social support. However, social support from friends remained the strongest predictor.

Wathen, C. N., MacGregor, J. C. D., Hammerton, J., Coben, J. H., Stewart, D. E., & MacMillan, H. L. (2012). [Priorities for research in child maltreatment, intimate partner violence and resilience to violence exposure: Results of an international Delphi consensus development process](#). *BMC Public Health*, 12(1), 684–696. DOI:10.1186/1471-2458-12-684

Intimate partner violence (IPV) and child maltreatment (CM) are major global public health problems. The Preventing Violence Across the Lifespan (PreVAil) Research Network, an international group of over 60 researchers and national and international knowledge-user partners in CM and IPV, sought to identify evidence-based research priorities in IPV and CM, with a focus on resilience, using a modified Delphi consensus development process. Review of existing empirical evidence, PreVAil documents and team discussion identified a starting list of 20 priorities in the following categories: resilience to violence exposure (RES), CM, and IPV, as well as priorities that cross-cut the content areas (CC), and others specific to research methodologies (RM) in violence research. PreVAil members (N = 47) completed two online survey rounds, and one round of discussions via three teleconference calls to rate, rank and refine research priorities. Research priorities were: to examine key elements of promising or successful programmes in RES/CM/IPV to build intervention pilot work; CC: to integrate violence questions into national and international surveys, and RM: to investigate methods for collecting and collating datasets to link data and to conduct pooled, meta and sub-group analyses to identify promising interventions for particular groups. These evidence-based research priorities, developed by an international team of violence, gender and mental health researchers and knowledge-user partners, are of relevance for prevention and resilience-oriented research in the areas of IPV and CM.

Wekerle, C., Waechter, R., & Chung, R. (2012). Contexts of vulnerability and resilience: Childhood maltreatment, cognitive functioning and close relationships. In M. Ungar (Ed.), *The social ecology of resilience: A handbook of theory and practice* (pp. 187–198). Springer. DOI:10.1007/978-1-4614-0586-3_16

Williams, J., & Nelson-Gardell, D. (2012). Predicting resilience in sexually abused adolescents. *Child Abuse & Neglect*, 36(1), 53–63. DOI:10.1016/j.chiabu.2011.07.004

This research examined factors that predicted resilience in sexually abused adolescents. Using Bronfenbrenner's Process–Person–Context–Time (PPCT) ecological model, this study considered the proximal and distal factors that would contribute to adolescents' reactions to sexual victimization. This correlational study used hierarchical regression analysis ($n = 237$) with cross-sectional data from the National Survey of Child and Adolescent Well-Being Wave I. This study found that school engagement, caregiver social support, hope and expectancy, caregiver education and SES predicted resilience. In line with the PPCT model, findings suggest that placing a greater emphasis on the contextual environment could improve support for adolescent resilience. Augmenting interventions that focus on individual change with those that address environmental factors may increase the benefits to adolescents affected by sexual abuse.

Zolkoski, S. M., & Bullock, L. M. (2012). Resilience in children and youth: A review. *Children and Youth Services Review*, 34(12), 2295–2303. DOI:10.1016/j.chilyouth.2012.08.009

Many children are reared in less than ideal family conditions (e.g., poverty, violence, substance abuse, family dissonance, family or personal illnesses). Situations such as these may inhibit the normal intellectual, social, and emotional development of children & youth, thus interfering with them reaching their full potential as adults. Conversely, many children encounter such adversities and fair well in spite of the challenges and may be considered to be resilient. This paper offers a review of the literature dating back to the 1970s to 2012. In addition, several monumental longitudinal studies dating back to the 1950s are included. The paper reviews the definition of resilience, origins and recent advances in researching resilience, protective factors, models of resilience, issues when researching resilience, measures of resilience, and resilience-based interventions.

Afifi, T. O., & MacMillan, H. L. (2011). Resilience following child maltreatment: A review of protective factors. *Canadian Journal of Psychiatry*, 56(5), 266-272.
DOI:10.1177/070674371105600505

Child maltreatment is linked to numerous adverse outcomes that can continue throughout the lifespan. However, impairment variability has been noted following child maltreatment, making it seem that some people are more resilient. Our review includes a discussion of how resilience is measured in child maltreatment research; a summary of the evidence for protective factors associated with resilience based on high quality studies; a discussion of how knowledge of protective factors can be applied to promote resilience among people exposed to child maltreatment; and finally, directions for future research. The databases MEDLINE & PsycINFO were searched for relevant citations up to July 2010 to identify key studies and evidence syntheses. Although comparability across studies is limited, family-level factors of stable family environment and supportive relationships appear to be consistently linked with resilience across studies. There was also evidence for some individual-level factors, such as personality traits, although proxies of intellect were not as strongly related to resilience following child maltreatment. Findings from resilience research needs to be applied to determine effective strategies and specific interventions to promote resilience & foster well-being among maltreated children.

Bogar, C. B., & Hulse-Killacky, D. (2011). Resiliency determinants and resiliency processes among female adult survivors of childhood sexual abuse. *Journal of Counseling & Development*, 84(3), 318-327. DOI:10.1002/j.1556-6678.2006.tb00411.x

This phenomenological, qualitative study examined resiliency determinants and resiliency formation among 10 women who had been sexually abused as children. An examination of the determinants and processes that facilitated resiliency in participants' adult lives revealed 5 determinant clusters (interpersonally skilled, competent, high self-

regard, spiritual, and helpful life circumstances) and 4 process clusters (coping strategies, refocusing & moving on, active healing, and achieving closure).

Masten, A. S. (2011). Resilience in children threatened by extreme adversity: Frameworks for research, practice, and translational synergy. *Development and Psychopathology*, 23(2), 493–506. DOI:10.1017/S0954579411000198

This article delineates parallel frameworks that grew out of the research on risk and resilience over the past 4 decades, a framework for research & a framework for practice, and then discusses the promise of an emerging synthesis. The research framework defined the meaning, models, and methods that successfully guided 4 waves of research to date on the nature & processes involved in human resilience. The applied framework emerged in response to urgent needs of children & families faced by adversity and those charged with helping them, resulting in guidelines for translating the unfolding but incomplete research evidence into action. The application of a resilience approach transformed practice in many fields concerned with promoting resilience in people at risk, revolutionizing the mission, models, measures, & methods of practice to align with the emphasis on positive adaptation and strengths defining a resilience-based approach but these interventions rarely translated back to inform and refine resilience theory in ways that would accelerate progress to promote resilience more effectively. The concluding section on translational synergy discusses the potential for a synthesis of basic & applied resilience frameworks as the next steps toward realizing the original objective and promise of resilience science.

Sapientza, J. K., & Masten, A. S. (2011). Understanding and promoting resilience in children and youth. *Current Opinion in Psychiatry*, 24(4), 267-273.
DOI:10.1097/YCO.0b013e32834776a8

In a context of global concern about the consequences of stress and extreme adversities, advances in theory and methods for studying human resilience have ushered in a new era of integrative, biopsychosocial research. This review highlights recent theory, findings, and implications of resilience research on young people. Resilience research has shifted toward dynamic system models with multiple levels of interaction, including research on the neurobiology of stress and adaption, epigenetic processes, and disasters. Growing evidence indicates individual differences in biological sensitivity to negative and positive experiences, including interventions. Early experiences show enduring programming effects on key adaptive systems, underscoring the importance of early intervention. Studies of developmental cascades demonstrate spreading effects of competence and symptoms over time, with important implications for the timing and targeting of interventions. Disaster research suggests guidelines for planning to protect children in the event of large-scale trauma. Research is integrating the study of resilience across system levels, with implications for promoting positive adaptation of young people faced with extreme adversity. However, studies on neurobiological and epigenetic processes are just beginning, and more research is needed on efficacy, as well as strategic timing and targeting, of interventions.

Ungar, M., & Liebenberg, L. (2011). Assessing resilience across cultures using mixed methods: Construction of the child and youth resilience measure. *Journal of Multiple Methods in Research*, 5(2), 126-149. DOI:10.1177/1558689811400607

An international team of investigators in 11 countries have worked collaboratively to develop a culturally and contextually relevant measure of youth resilience, the Child and Youth Resilience Measure (CYRM-28). The team used a mixed methods design that facilitated understanding of both common and unique aspects of resilience across

cultures. Quantitative and qualitative stages to its development ensure the CYRM-28 has good content-related validity across research sites. Crossover comparison analyses of the findings from the quantitative administration of the pilot measure with 1,451 youth and qualitative interviews with 89 youth support the CYRM-28 as a culturally sensitive measure of youth resilience. The implications of this mixed methods approach to the development of measures for cross-cultural research are discussed.

Windle, G., Bennett, K. M., & Noyes, J. (2011). [A methodological review of resilience measurement scales](#). *Health and Quality of Life Outcomes*, 9(8), 1-18.
DOI:10.1186/1477-7525-9-8

The evaluation of interventions and policies designed to promote resilience, and research to understand the determinants and associations, require reliable and valid measures to ensure data quality. This paper systematically reviews the psychometric rigour of resilience measurement scales developed for use in general and clinical populations. Eight electronic abstract databases and the internet were searched and reference lists of all identified papers were hand searched. The focus was to identify peer reviewed journal articles where resilience was a key focus and/or is assessed. Two authors independently extracted data and performed a quality assessment of the scale psychometric properties. Nineteen resilience measures were reviewed; four of these were refinements of the original measure. All the measures had some missing information regarding the psychometric properties. Overall, the Connor-Davidson Resilience Scale, the Resilience Scale for Adults and the Brief Resilience Scale received the best psychometric ratings. The conceptual and theoretical adequacy of a number of the scales was questionable. We found no current 'gold standard' amongst 15 measures of resilience. A number of the scales are in the early stages of development, and all require further validation work. Given increasing interest in resilience from major international

funders, key policy makers and practice, researchers are urged to report relevant validation statistics when using the measures.

Davidson, G., Devaney, J., & Spratt, T. (2010). The impact of adversity in childhood on outcomes in adulthood research lessons and limitations. *Journal of Social Work, 10*(4), 369–390. DOI:10.1177/1468017310378783

Current UK Government policy is concerned with the possible connections between childhood adversity, social exclusion and negative outcomes in adulthood. Understanding the impact of adverse childhood experiences on outcomes in adulthood is therefore key to informing effective policy and practice. In this article, the research on the impact of childhood adversity on outcomes in adulthood is reviewed in the broad categories of: mental health and social functioning; physical health; offending; service use; and economic impact. The literature on resilience that focuses on those who experience adversity, but do not have associated negative outcomes is also briefly considered. The strengths and limitations of the range of research methods used are then examined. Previous studies have tended to focus on specific forms of adversity, predominantly abuse and neglect, and either: specific populations and specific outcomes; specific populations and general outcomes; or general populations and specific outcomes. This means there may be incomplete understanding of the inputs (the range of adverse experiences in childhood), the processes (how these may affect people) and the outcomes (across domains in adulthood). It is concluded that it is important for social work researchers to engage in the current debate about how to prevent harmful childhood adversity and there is an important gap in the research for more interdisciplinary large-scale general population studies that consider the full range of childhood adversity and associated impacts across time and the possible processes involved.

Atkinson, P. A., Martin, C. R., & Rankin, J. (2009). Resilience revisited. *Journal of Psychiatric and Mental Health Nursing*, 16(2), 137–145. DOI:10.1111/j.1365-2850.2008.01341.x

Contemporary reports of individuals experiencing severe adversity have led to renewed debate regarding the processes that mediate a person's ability to withstand extremes of both psychological and physical abuse. The capacity to recover from extremes of trauma and stress is termed resilience. This important concept is of key relevance to health professionals as resilience also defines a more general concept of responding to challenges that affect the individual in terms of their health behaviour. The evidence base in this area is noteworthy in that much research is based on historical case studies that, although rich in both content and appeal, also fall short of contemporary rigour in terms of validity. This review will describe the evidence for the salience of the concept of the resilience to health and mental health and the implications of the domain to the healthcare professional.

Martinez-Torteya, C., Bogat, G. A., von Eye, A., & Levendosky, A. A. (2009). Resilience among children exposed to domestic violence: The role of risk and protective factors. *Child Development*, 80(2), 562–577. DOI:10.1111/j.1467-8624.2009.01279.x

Individual and family characteristics that predict resilience among children exposed to domestic violence (DV) were examined. Mother–child dyads ($n = 190$) were assessed when the children were 2, 3, and 4 years of age. DV-exposed children were 3.7 times more likely than nonexposed children to develop internalizing or externalizing problems. However, 54% of DV-exposed children maintained positive adaptation and were characterized by easy temperament (odds ratio [OR] = .39, $d = .52$) and nondepressed mothers (OR = 1.14, $d = .07$), as compared to their nonresilient counterparts. Chronic DV was associated with maternal depression, difficult child temperament, and internalizing or externalizing symptoms. Results underscore heterogeneous outcomes among DV-

exposed children and the influence of individual and family characteristics on children's adaptation.

American Psychological Association Task Force on Resilience and Strengths in Black Children and Adolescents. (2008). [Resilience in African American children and adolescents: A vision for optimal development](https://www.apa.org/pi/families/resources/task-force/resilience-af-am).
<https://www.apa.org/pi/families/resources/task-force/resilience-af-am>

McClure, F. H., Chavez, D. V., Agars, M. D., Peacock, M. J., & Matosian, A. (2008). Resilience in sexually abused women: Risk and protective factors. *Journal of Family Violence*, 23(2), 81-88. DOI:10.1007/s10896-007-9129-4

The goal of this study was to evaluate the contribution of family characteristics (family conflict and cohesion) and abuse characteristics (age at which abuse occurred, abuse severity, and relationship to perpetrator) to resilience (self-acceptance, ability to engage in positive relationships with others, and environmental mastery) in a sample of one hundred and 77 university women who had experienced childhood sexual abuse (CSA). The respondents completed a questionnaire that focused on childhood experiences including family environment in childhood, characteristics of the abuse they had experienced in addition to degree of current resilience defined as their level of well-being (self-acceptance, ability to engage in positive relationships with others, and environmental mastery). The results indicated that family characteristics accounted for 13–22% of the variance in the well-being outcomes. In contrast, abuse characteristics accounted for 3% or less of the variance in the well-being outcomes. The clinical implications of these findings are discussed.

Suzuki, S. L., Geffner, R., & Bucky, S. F. (2008). The experiences of adults exposed to intimate partner violence as children: An exploratory qualitative study of resilience and protective factors. *Journal of Emotional Abuse*, 8(1-2), 103-121. DOI:10.1080/10926790801984523

Research indicates a large percentage of children exposed to intimate partner violence (IPV) experience psychological, social, cognitive, behavioral, and physical problems. These deficits can have overwhelming long-term problems for children. Although there is some evidence of the intergenerational cycle of violence, not all children exposed to IPV encounter abuse in their adult relationships. This study qualitatively explored the protective factors or resiliencies that contribute to adaptive outcomes of adults exposed to IPV as children. Ten major and five minor themes emerged from this analysis. These themes were organized into internal factors/individual characteristics, family factors, and external factors. Implications for practice are also discussed.

Breno, A. L., & Galupo, M. P. (2007). Sexual abuse histories of young women in the US child welfare system: A focus on trauma-related beliefs and resilience. *Journal of Child Sexual Abuse*, 16(2), 97-113. DOI:10.1300/J070v16n02_06

This research provides descriptive data regarding sexual abuse histories of high-functioning women ($N = 84$; 18-25 years old) previously in the child welfare system. Placement histories of foster youth who were sexually abused were distinct. Girls with a history of sexual abuse were more likely to have been in restrictive housing and changed placements twice as often as girls with no history of sexual abuse. Trauma-related beliefs (TRB) subsequent to sexual abuse varied depending upon where sexual abuse occurred. TRB scores were negatively correlated with resilience and positively correlated with number of sexual abusers. Powerlessness was found to make a significant contribution to resiliency scores above and beyond foster care and abuse demographics.

Daigneault, I., Hébert, M., & Tourigny, M. (2007). Personal and interpersonal characteristics related to resilient developmental pathways of sexually abused adolescents. *Child and Adolescent Psychiatric Clinics of North America*, 16(2), 415–434. DOI:10.1016/j.chc.2006.11.002

Childhood sexual abuse is a prevalent and important social problem. In this article the authors' main objective is the exploration of resilience among sexually abused adolescents under child protection services care in Quebec, Canada. The introduction provides an overview of the impact of childhood sexual abuse and conceptualizations of trauma, resilience, recovery, and factors linked to resilience profiles. The remainder of the article discusses a study of resilience and factors predicting resilience over a 5-month period for a sample of 86 teenage girls under child protection services care.

DuMont, K. A., Widom, C. S., & Czaja, S. J. (2007). Predictors of resilience in abused and neglected children grown-up: The role of individual and neighborhood characteristics. *Child Abuse & Neglect*, 31(3), 255–274. DOI:10.1016/j.chiabu.2005.11.015

This paper examines individual, family, and neighborhood level predictors of resilience in adolescence and young adulthood and describes changes in resilience over time from adolescence to young adulthood in abused and neglected children grown up. We use documented cases of childhood physical and sexual abuse and neglect ($n = 676$) from a Midwestern county area during the years 1967–1971 and information from official records, census data, psychiatric assessments, and self-reports obtained through 1995. Analyses involve logistic regressions, replicated with Mplus to test for possible contextual effects. Almost half (48%) of the abused and neglected children in adolescence and nearly one-third in young adulthood were resilient. Over half of those who were resilient in adolescence remained resilient in young adulthood, whereas 11% of the non-resilient adolescents were resilient in young adulthood. Females were more likely to be resilient during both time periods. Being white, non-Hispanic decreased and growing up in a

stable living situation increased the likelihood of resilience in adolescence, but not in young adulthood. Stressful life events and a supportive partner promoted resilience in young adulthood. Neighborhood advantage did not exert a direct effect on resilience, but moderated the relationship between household stability and resilience in adolescence and between cognitive ability and resilience in young adulthood.

Walsh, K., Blaustein, M., Knight, W. G., Spinazzola, J., & van der Kolk, B. A. (2007). Resiliency factors in the relation between childhood sexual abuse and adulthood sexual assault in college-age women. *Journal of Child Sexual Abuse, 16*(1), 1-17.
DOI:10.1300/J070v16n01_01

Research has suggested that childhood sexual abuse (CSA) may be a risk factor for adulthood sexual assault. This study examined associations between CSA experiences, cognitive resiliency variables, and revictimization. Participants were 73 college-age females who completed self-report questionnaires assessing CSA, adult assault, self-efficacy, locus of control (LOC), and coping styles. Sexual assault was categorized as forced or coerced assault based on the tactics used by the perpetrator. Results indicated that CSA alone was the strongest independent predictor of forced adult assault; however, LOC and positive coping were associated with resiliency to coercive sexual assault. The current findings have clinical implications in that LOC and coping styles are characteristics that can be enhanced through therapy.

Ahern, N. R., Kiehl, E. M., Sole, M., & Byers, J. (2006). A review of instruments measuring resilience. *Issues in Comprehensive Pediatric Nursing, 29*(2), 103-125.
DOI:10.1080/01460860600677643

The objectives of the study were to evaluate the psychometric properties and appropriateness of instruments for the study of resilience in adolescents. A search was completed using the terms resilience and instruments or scales using the EBSCO

database (CINAHL, PreCINAHL, and Academic Search Premier), MEDLINE, PsychINFO and PsychARTICLES, and the Internet. After instruments were identified, a second search was performed for studies reporting the psychometric development of these instruments. Using inclusion and exclusion criteria, six psychometric development of instrument studies were selected for a full review. A data extraction table was used to compare the six instruments. Two of the six instruments (Baruth Protective Factors Inventory [BPFI] and Brief-Resilient Coping Scale) lacked evidence that they were appropriate for administration with the adolescent population due to lack of research applications. Three instruments (Adolescent Resilience Scale [ARS], Connor-Davidson Resilience Scale, and Resilience Scale for Adults) had acceptable credibility but needed further study in adolescents. One instrument (Resilience Scale [RS]) was determined to be the best instrument to study resilience in the adolescent population due to psychometric properties of the instrument and applications in a variety of age groups, including adolescence. Findings of this review indicate that the RS is the most appropriate instrument to study resilience in the adolescent population. While other instruments have potential (e.g., ARS, BPFI) as they were tested in the adolescent and young adult populations, they lack evidence for their use at this time. An evaluation of the review and recommendations are discussed.

Edmond, T., Auslander, W., Elze, D., & Bowland, S. (2006). Signs of resilience in sexually abused adolescent girls in the foster care system. *Journal of Child Sexual Abuse, 15*(1), 1-29. DOI:10.1300/J070v15n01_01

Childhood sexual abuse has been shown to result in numerous negative emotional and psychological difficulties that can last a lifetime (e.g., Beitchman, Zucker, Hood, DaCosta, & Cassavia, 1992; Desai, Arias, Thompson, & Basile, 2002; Thompson, Arias, Basile, & Desai, 2002). For those sexual abuse survivors who are placed in the foster care system, there is the hope of the abuse ending but it comes at the cost of separation from their families.

Both events constitute significant stressors that can disrupt normal development. Youths who have had these experiences are psychologically at risk for developing mental health and behavioral problems that can negatively affect their life trajectories. Yet, there are those who, despite having had such painful experiences as child sexual abuse and placement in the foster care system, appear to be functioning well. This paper provides an empirical examination of a group of sexually abused adolescent girls who are in the foster care system, some of whom seem to be psychologically well adjusted and some of whom are struggling with serious mental health and behavioral problems. This within group difference provides an opportunity to study signs of resilience in a population that has not been previously studied by resilience researchers.

Greenberg, M. T. (2006). Promoting resilience in children and youth: Preventive interventions and their interface with neuroscience. *Annals of the New York Academy of Sciences*, 1094(1), 139–150. DOI:10.1196/annals.1376.013

Preventive interventions focus on reducing risk and promoting protective factors in the child as well as their cultural ecologies (family, classroom, school, peer groups, neighborhood, etc). By improving competencies in both the child and their contexts many of these interventions promote resilience. Although there are now a substantial number of preventive interventions that reduce problem behaviors and build competencies across childhood and adolescence, there has been little integration with recent findings in neuropsychology and neuroscience. This article focuses on the integration of prevention research and neuroscience in the context of interventions that promote resilience by improving the executive functions (EF); inhibitory control, planning, and problem solving skills, emotional regulation, and attentional capacities of children and youth. Illustrations are drawn from recent randomized controlled trials of the Promoting Alternative Thinking Strategies (PATHS) curriculum. The discussion focuses on the next steps in transdisciplinary research in prevention and social neuroscience.

Masten, A. S., & Shaffer, A. (2006). How families matter in child development: Reflections from research on risk and resilience. In A. Clarke-Stewart & J. Dunn (Eds.), *Families count: Effects on child and adolescent development* (pp. 5–25). Cambridge University Press. DOI:10.1017/CBO9780511616259.002

Alvord, M. K., & Grados, J. J. (2005). Enhancing resilience in children: A proactive approach. *Professional Psychology: Research and Practice*, 36(3), 238–245. DOI:10.1037/0735-7028.36.3.238

Many clinical practitioners today are interested in helping children be more resilient. The authors briefly review the literature and identify protective factors that are related to or foster resilience in children. After discussing individual and family intervention strategies currently in use, the authors present a practical, proactive, resilience-based model that clinicians may use in a group intervention setting. The model entails interactive identification of protective factors with children, free play and behavioral rehearsal, training in relaxation and self-control techniques, practice in generalizing skills acquired, and active parent involvement. Implications of this group intervention model are discussed.

Boyden, J., & Mann, G. (2005). Children's risk, resilience, and coping in extreme situations. In Author (Ed.), *Handbook for working with children and youth: Pathways to resilience across cultures and contexts* (pp. 3–26). Sage Publications, Inc. DOI:10.4135/9781412976312

Fergus, S., & Zimmerman, M. A. (2005). Adolescent resilience: A framework for understanding healthy development in the face of risk. *Annual Review of Public Health*, 26, 399–419. DOI:10.1146/annurev.publhealth.26.021304.144357

Adolescent resilience research differs from risk research by focusing on the assets and resources that enable some adolescents to overcome the negative effects of risk

exposure. We discuss three models of resilience—the compensatory, protective, and challenge models—and describe how resilience differs from related concepts. We describe issues and limitations related to resilience and provide an overview of recent resilience research related to adolescent substance use, violent behavior, and sexual risk behavior. We then discuss implications that resilience research has for intervention and describe some resilience-based interventions.

Flores, E., Cicchetti, D., & Rogosch, F. A. (2005). Predictors of resilience in maltreated and nonmaltreated Latino children. *Developmental Psychology*, 41(2), 338–351.
DOI:10.1037/0012-1649.41.2.338

To date, few studies have sought to investigate the effects of child maltreatment and processes influencing maladaptation and resilience in Latino children. In the current investigation, multiple aspects of functioning, personal resources, and relationship features were examined in school-age maltreated and nonmaltreated Latino children. Maltreated Latino children were found to have fewer areas of resilient functioning. Ego-resiliency and ego-control, as personal resources, and the ability to form a positive relationship with an adult figure outside of the immediate family predicted resilience. However, certain aspects of interpersonal functioning were differentially related to resilience for maltreated and nonmaltreated Latino children. These findings have implications for understanding how resilience can be promoted in maltreated and nonmaltreated Latino children.

Tedeschi, R. G., & Kilmer, R. P. (2005). [Assessing strengths, resilience, and growth to guide clinical interventions](#). *Professional Psychology: Research and Practice*, 36(3), 230–237. DOI:10.1037/0735-7028.36.3.230

Recently, the field of mental health has incorporated a growing interest in strengths, resilience, and growth, psychological phenomena that may be associated with healthy

adjustment trajectories and profitably integrated into strategies for clinical assessment and practice. This movement constitutes a significant shift from traditional deficit-oriented approaches. Addressing clinical practitioners, this article (a) provides a broad overview of these constructs and phenomena, (b) discusses their relevance for clinical assessment and intervention, and (c) describes selected strategies and approaches for conducting assessments that can guide intervention.

Bonanno, G. A. (2004). [Loss, trauma, and human resilience: Have we underestimated the human capacity to thrive after extremely aversive events?](#). *American Psychologist*, 59(1), 20–28. DOI:10.1037/0003-066X.59.1.20

Many people are exposed to loss or potentially traumatic events at some point in their lives, and yet they continue to have positive emotional experiences and show only minor and transient disruptions in their ability to function. Unfortunately, because much of psychology's knowledge about how adults cope with loss or trauma has come from individuals who sought treatment or exhibited great distress, loss and trauma theorists have often viewed this type of resilience as either rare or pathological. The author challenges these assumptions by reviewing evidence that resilience represents a distinct trajectory from the process of recovery, that resilience in the face of loss or potential trauma is more common than is often believed, and that there are multiple and sometimes unexpected pathways to resilience.

Masten, A. S. (2004). Regulatory processes, risk, and resilience in adolescent development. *Annals of the New York Academy of Sciences*, 1021(1), 310–319. DOI:10.1196/annals.1308.036

This article highlights the potential of developmental psychopathology as a useful integrative perspective for the challenging task of linking the study of brain development and adolescent behavior in context; it considers clues from behavioral research on

resilience on the nature of regulatory processes as risks and assets, or vulnerabilities and protective factors for the development of competence and psychopathology in adolescence; and it advocates more integrative neurobehavioral research on risk and resilience in adolescent development.

Trickett, P. K., Kurtz, D. A., & Pizzigati, K. (2004). Resilient outcomes in abused and neglected children: Bases for strengths-based intervention and prevention policies. In K. I. Maton, C. J. Schellenbach, B. J. Leadbeater, & A. L. Solarz (Eds.), *Investing in children, youth, families, and communities: Strengths-based research and policy* (pp. 73–95). American Psychological Association.
DOI:10.1037/10660-005

Wilcox, D. T., Richards, F., & O'Keeffe, Z. C. (2004). Resilience and risk factors associated with experiencing childhood sexual abuse. *Child Abuse Review*, 13(5), 338–352.
DOI:10.1002/car.862

The effects of child sexual abuse are wide-ranging and impact on children, families and health/social care systems. The authors review this literature, examining the shortcomings of the 'victim-offender' model, and consider the complex, multifactorial nature of this question. Factors associated with a progression from victim to perpetrator are explored and the prevalence of abuse in the general population is also discussed. Protective as well as risk factors are considered and the pivotal role of 'personal reliance' is considered as it relates to empowering damaged young people to become healthy adults. The authors also discuss implications for the treatment of young people who become abusers. Two case examples are briefly reported upon and the article concludes with a consideration of ways to helpfully address the needs of children who become sexual perpetrators.

Bolger, K. E., & Patterson, C. J. (2003). Sequelae of child maltreatment: Vulnerability and resilience. In S. S. Luthar (Ed.), *Resilience and vulnerability: Adaptations in the context of childhood adversities* (pp. 156–181). Cambridge University Press. DOI:10.1017/CBO9780511615788.009

Reynolds, A. J., & Ou, S. (2003). Promoting resilience through early childhood intervention. In S. S. Luthar (Eds.), *Resilience and vulnerability: Adaptation in the context of childhood adversities*. Cambridge University Press. DOI:10.1017/CBO9780511615788

Banyard, V. L., Williams, L. M., Siegel, J. A., & West, C. M. (2002). Childhood sexual abuse in the lives of black women: Risk and resilience in a longitudinal study. *Women & Therapy*, 25(3–4), 45–58. DOI:10.1300/J015v25n03_04

Childhood sexual abuse (CSA) is associated with long-term mental health consequences. This article reviews the results of one longitudinal study, whose sample consisted primarily of African American women. The purpose is to give voice to an understudied group of CSA survivors and to highlight the variability in risk and protective factors. Key findings related to mental health consequences, re-traumatization, and resilience are reviewed and set within the broader context of research on African American women and child sexual abuse. Implications for future research and clinical practice are discussed.

Lambie, I., Seymour, F., Lee, A., & Adams, P. (2002). Resiliency in the victim–offender cycle in male sexual abuse. *Sexual Abuse*, 14(1), 31–48. DOI:10.1177/107906320201400103

The prevalence of childhood sexual abuse in child molesters is considerably higher than that in the general population. This finding had led to the “victim–offender cycle” being popularized as an explanation for sexual offending. However, not all child molesters were victimized as children, so it is of interest to examine the factors that contribute to the victim–offender cycle or, conversely, resiliency. This study examined the “moderating

factors” that may prevent a male victim of sexual abuse from entering the victim–offender cycle. Two groups were interviewed as part of the study: a “resilient group” ($n = 47$) and a victim–offender group ($n = 41$). After correction for age and education level, the resilient group were less likely to have fantasized and masturbated about the abuse, less likely to report deriving pleasure from the abuse, more likely to have had frequent social contact with adolescent peers and to have had more family and nonfamily support during childhood. The findings support the need for multifactorial models of resiliency, the victim–offender cycle, and sexual offending. Recommendations about the prevention of the victim–offender cycle are made, including the need for a thorough systemic assessment of all male victims of sexual abuse and the involvement of their family system in counseling.

McGloin, J. M., & Widom, C. S. (2001). Resilience among abused and neglected children grown up. *Development and Psychopathology*, 13(4), 1021–1038.
DOI:10.1017/s095457940100414x

Although an extensive literature has accumulated documenting the maladaptive outcomes associated with childhood victimization, a limited body of knowledge addresses resilience. This paper sought to operationalize the construct of resilience across a number of domains of functioning and time periods and to determine the extent to which abused and neglected children grown up demonstrate resilience. Substantiated cases of child abuse and neglect from 1967 to 1971 were matched on gender, age, race, and approximate family social class with non-abused and non-neglected children and followed prospectively into young adulthood. Between 1989 and 1995, 1,196 participants (676 abused and neglected and 520 controls) were administered a 2-hr in-person interview, including a psychiatric assessment. Resilience requires meeting the criteria for success across six of eight domains of functioning: employment, homelessness, education, social activity, psychiatric disorder, substance abuse, and two domains

assessing criminal behavior (official arrest and self-reports of violence). Results indicate that 22% of abused and neglected individuals meet the criteria for resilience. More females met the criteria for resilience and females were successful across a greater number of domains than males. We speculate on the meaning of these findings and discuss implications for the child maltreatment field. Limitations of the study are also acknowledged.

Werner, E. E., & Smith, R. S. (2001). *Journeys from childhood to mid-life: Risk, resilience, and recovery*. Cornell University Press.

Luthar, S. S., & Cicchetti, D. (2000). [The construct of resilience: Implications for interventions and social policies](#). *Development and Psychopathology*, 12(4), 857–885. DOI:10.1017/s0954579400004156

The focus of this article is on the interface between research on resilience—a construct representing positive adaptation despite adversity—and the applications of this work to the development of interventions and social policies. Salient defining features of research on resilience are delineated, as are various advantages, limitations, and precautions linked with the application of the resilience framework to developing interventions. For future applied efforts within this tradition, a series of guiding principles are presented along with exemplars of existing programs based on the resilience paradigm. The article concludes with discussions of directions for future work in this area, with emphases on an enhanced interface between science and practice, and a broadened scope of resilience-based interventions in terms of the types of populations, and the types of adjustment domains, that are encompassed.

Luthar, S. S., Cicchetti, D., & Becker, B. (2000). [The construct of resilience: A critical evaluation and guidelines for future work](#). *Child Development*, 71(3), 543–562.
DOI:10.1111/1467-8624.00164

This paper presents a critical appraisal of resilience, a construct connoting the maintenance of positive adaptation by individuals despite experiences of significant adversity. As empirical research on resilience has burgeoned in recent years, criticisms have been levied at work in this area. These critiques have generally focused on ambiguities in definitions and central terminology; heterogeneity in risks experienced and competence achieved by individuals viewed as resilient; instability of the phenomenon of resilience; and concerns regarding the usefulness of resilience as a theoretical construct. We address each identified criticism in turn, proposing solutions for those we view as legitimate and clarifying misunderstandings surrounding those we believe to be less valid. We conclude that work on resilience possesses substantial potential for augmenting the understanding of processes affecting at-risk individuals. Realization of the potential embodied by this construct, however, will remain constrained without continued scientific attention to some of the serious conceptual and methodological pitfalls that have been noted by skeptics and proponents alike.

Werner, E. (2000). Protective factors and individual resilience. In J. Shonkoff, & S. Meisels (Eds.), *Handbook of early childhood intervention* (2nd ed., pp. 115–132). Cambridge University Press. DOI:10.1017/CBO9780511529320.008

Wyman, P. A., Sandler, I., Wolchik, S., & Nelson, K. (2000). Resilience as cumulative competence promotion and stress protection: Theory and intervention. In D. Cicchetti, J. Rappaport, I. Sandler, & R. P. Weissberg (Eds.), *The promotion of wellness in children and adolescents* (pp.133–184). Child Welfare League of America Press. DOI:10.1007/0-387-23824-7_2

Heller, S. S., Larrieu, J. A., D'Imperio, R., & Boris, N. W. (1999). Research on resilience to child maltreatment: Empirical considerations. *Child Abuse & Neglect*, 23(4), 321-338. DOI:10.1016/s0145-2134(99)00007-1

To review the current research literature on resilience to maltreatment in childhood and adolescence. First, this paper addresses the methodological issues applicable to this area of research. Second, it reviews the empirical literature on the protective factors believed to contribute to resilience to maltreatment and offers commentary on specific issues raised in each study. The current findings are summarized and an integrated descriptive picture, based on the extant literature of the processes and factors contributing to the development of resilience to childhood maltreatment is outlined. Based on this extensive review and critique of the current empirical literature on resilience to maltreatment, suggestions for future investigations in this area are recommended.

Masten, A. S., Hubbard, J. J., Gest, S. D., Tellegen, A., Garmezy, N., & Ramirez, M. L. (1999). Competence in the context of adversity: Pathways to resilience and maladaptation from childhood to late adolescence. *Development and Psychopathology*, 11(1), 143-169. DOI:10.1017/s0954579499001996

Competent outcomes in late adolescence were examined in relation to adversity over time, antecedent competence and psychosocial resources, in order to investigate the phenomenon of resilience. An urban community sample of 205 (114 females, 90 males; 27% minority) children were recruited in elementary school and followed over 10 years. Multiple methods and informants were utilized to assess three major domains of competence from childhood through adolescence (academic achievement, conduct, and peer social competence), multiple aspects of adversity, and major psychosocial resources. Both variable-centered and person-centered analyses were conducted to test the hypothesized significance of resources for resilience. Better intellectual functioning and parenting resources were associated with good outcomes across competence domains, even in the context of severe, chronic adversity. IQ and parenting appeared to

have a specific protective role with respect to antisocial behavior. Resilient adolescents (high adversity, adequate competence across three domains) had much in common with their low-adversity competent peers, including average or better IQ, parenting, and psychological well-being. Resilient individuals differed markedly from their high adversity, maladaptive peers who had few resources and high negative emotionality. Results suggest that IQ and parenting scores are markers of fundamental adaptational systems that protect child development in the context of severe adversity.

Cicchetti, D., & Rogosch, F. A. (1997). The role of self-organization in the promotion of resilience in maltreated children. *Development and Psychopathology*, 9(4), 797-815. DOI:10.1017/s0954579497001442

The dynamic nature of resilience necessitates that children from high-risk backgrounds who are functioning adaptively despite experiences of adversity must be examined over time. In the current investigation, the adaptation of school-age maltreated and non-maltreated socioeconomically disadvantaged children was examined over 3 consecutive years. In accord with predictions, a higher percentage of non-maltreated children than of maltreated children were found to be resilient. Moreover, a higher percentage of maltreated than of non-maltreated children were shown to exhibit functioning consistently in the low adaptive range. Differential predictors of resilience were found in maltreated and non-maltreated children. Specifically, for maltreated children, positive self-esteem, ego resilience, and ego overcontrol predicted resilient functioning, whereas relationship features were more influential for non-maltreated children. These findings are discussed in relation to the unfolding of resilient self-organizational strivings in maltreated and non-maltreated children.

Himelein, M. J., & McElrath, J. A. V. (1996). Resilient child sexual abuse survivors: Cognitive coping and illusion. *Child Abuse & Neglect*, 20(8), 747–758. DOI:10.1016/0145-2134(96)00062-2

This research examined cognitive coping strategies associated with resilience in a nonclinical sample of child sexual abuse (CSA) survivors. In Study 1, 180 college women completed surveys assessing self-enhancing cognitive distortions of reality, known as positive illusions, and CSA history. CSA survivors and nonvictimized women were found to be equally likely to engage in illusion, and for both groups, measures of illusion were strongly associated with psychological well-being. In Study 2, a qualitative study, a subsample of 20 CSA survivors from Study 1 were interviewed regarding their efforts to cope with CSA. Analysis was focused on comparisons between well-adjusted and poorly-adjusted women. The high adjustment group revealed a greater tendency to engage in four types of cognitive strategies: disclosing and discussing CSA, minimization, positive reframing, and refusing to dwell on the experience. The results of both studies highlight the importance of cognitive reappraisal in CSA recovery. Implications for therapists working with CSA survivors are discussed.

Rak, C. F., & Patterson, L. E. (1996). Promoting resilience in at-risk children. *Journal of Counseling and Development*, 74(4), 368–373.
DOI:10.1002/j.15566676.1996.tb01881.x

Intervening effectively with youths at risk from early deprivation, family dysfunction, poverty, abuse, and other factors is a major concern for educational and social service policymakers. Current research suggests that a majority of at-risk youths do not experience drastic outcomes, but many exhibit protective factors that buffer them from negative consequences. Longitudinal studies from Hawaii, the continental United States, and Great Britain have identified several personality, familial, and environmental variables that promote resiliency in youths at risk. This article discusses these variables

and provides counselors with an assessment technique and strategies to promote a salutogenesis perspective.

Spaccarelli, S., & Kim, S. (1995). Resilience criteria and factors associated with resilience in sexually abused girls. *Child Abuse & Neglect*, 19(9), 1171-1182.
DOI:10.1016/01452134(95)00077-1

Alternative measures of resilience and correlates of resilience were examined in a sample of 43 sexually abused girls who were assessed using a self-administered interview at the time of intake for psychotherapy. Results indicated relatively high levels of disagreement as to which girls were resilient using maintenance of social competence and absence of clinical levels of symptomatology as alternative criteria. Most girls that had maintained age-normative levels of social competence were, nonetheless, manifesting clinically significant levels of symptoms. A warm and supportive relationship with a nonoffending parent was a strong correlate of resilience, regardless of which criteria was used. Lower levels of abuse related stress, fewer negative cognitive appraisals of the abusive relationship, and less reliance on aggressive coping behaviors were also significant predictors of resilience based on the absence of clinical levels of symptomatology. However, parental support and level of abuse stress were the only two variables to enter a logistic regression model predicting resilience. The research and clinical implications of these findings are discussed.

Herrenkohl, E. C., Herrenkohl, R. C., & Egolf, B. (1994). Resilient early school-age children from maltreating homes: Outcomes in late adolescence. *American Journal of Orthopsychiatry*, 64(2), 301-309. DOI:10.1037/h0079517

The interaction of individual and environmental characteristics over time in children from maltreating families is examined in order to explore the origins, scope, and stability of resilience in children from abusive and neglectful home environments. Response to

supportive influences in the extended family and wider community, as well as determination to be different from abusive parents, is emphasized as crucial to resilient behavior and is illustrated by a case study.

Cicchetti, D., Rogosch, F., Lynch, M., & Holt, K. (1993). Resilience in maltreated children: Process leading to adaptive children. *Development and Psychopathology*, 5(4), 629–647. DOI:10.1017/S0954579400006209

Evidence for resilience, competent functioning despite severe adversity, was investigated in school-age, disadvantaged maltreated ($N = 127$) and nonmaltreated ($N = 79$) children attending a summer camp program. Multiple areas of adaptation (social adjustment, risk for school difficulty, psychopathology) were assessed from self, peer, and camp counselor perspectives and school records. A composite index of adaptive functioning was developed, and levels of competence were delineated. Personality dimensions and personal resources, including cognitive maturity, self-esteem, ego-resiliency, and ego-control, were evaluated as mechanisms promoting individual differences in successful adaptation. Maltreated children as a group evidenced lower overall competence when compared to nonmaltreated children. An equal proportion of maltreated and nonmaltreated children, however, demonstrated high levels of competence, whereas more maltreated children than nonmaltreated children evidenced low levels of competence. Ego-resiliency, ego-control, and self-esteem were each found to predict individual differences in competent functioning. Evidence for the differential role of ego-control in promoting competence for maltreated versus nonmaltreated children was found. The results are discussed in terms of mechanisms contributing to resilient outcomes in maltreated children and the implications of the study of resilience for the field of developmental psychopathology.

Valentine, L., & Feinauer, L. L. (1993). Resilience factors associated with female survivors of childhood sexual abuse. *American Journal of Family Therapy*, 21(3), 216–224. DOI:10.1080/01926189308250920

The 22 women interviewed in this study were in a high-risk group for low self-esteem, depression, shame, and other long-term interpersonal difficulties due to their survival of childhood sexual abuse. Unlike many of their contemporaries, they have been able to have relationships, stable careers, and healthy personalities. This paper explores the variables and patterns gathered from their interviews. The resiliency themes extracted included: the ability to find emotional support outside the family; self-regard or the ability to think well of oneself; spirituality; external attribution of blame and cognitive style; and inner-directed locus of control.

Garnezy, N. (1991). Resilience in children's adaptation to negative life events and stressed environments. *Pediatric Annals*, 20(9), 459–460. DOI:10.3928/0090-4481-19910901-05

Functional adequacy (the maintenance of competent functioning despite an interfering emotionality) is a benchmark of resilient behavior under stress. While resilient adults can be identified as adults who once experienced a great deal of despair as children, I am not prepared to mark off the construct of resilience because such people may carry with them a realistic baggage of sadness and unhappiness. The very nature of despair that is present for children of the ghetto, the status of minority children in today's America, all have a reality that can neither be ignored nor denied. Perhaps a portion of resilient behavior is the evaluative awareness of a difficult reality combined with a commitment to struggle, to conquer the obstacle, and to achieve one's goals despite the negative circumstances to which one has been exposed, which were and remain evocative of sadness.

Luthar, S. S. (1991). [Vulnerability and resilience: A study of high-risk adolescents](#). *Child Development*, 62(3), 600–616. DOI:10.1111/j.1467-8624.1991.tb01555.x

Factors that allow children to maintain socially competent behaviors despite stress were examined among 144 inner-city ninth-grade students with a mean age of 15.3 years. Stress was operationalized by scores on a negative life events scale, and definitions of social competence were based on peer ratings, teacher ratings, and school grades. Moderator variables examined included intelligence, internal locus of control, social skills, ego development, and positive life events. Following theoretical models by Garmezy and Rutter, distinctions were made between compensatory factors (which are directly related to competence) and protective/vulnerability factors (which interact with stress in influencing competence). Ego development was found to be compensatory against stress. Internality and social skills proved to be protective factors, while intelligence and positive events were involved in vulnerability processes. This study also revealed that children labeled as resilient were significantly more depressed and anxious than were competent children from low stress backgrounds.

Masten, A. S., Best, K. M., & Garmezy, N. (1990). Resilience and development: Contributions from the study of children who overcome adversity. *Development and Psychopathology*, 2(4), 425–444. DOI:10.1017/S0954579400005812

This article reviews the research on resilience in order to delineate its significance and potential for understanding normal development. Resilience refers to the process of, capacity for, or outcome of successful adaptation despite challenging or threatening circumstances. Three resilience phenomena are reviewed: (a) good outcomes in high-risk children, (b) sustained competence in children under stress, and (c) recovery from trauma. It is concluded that human psychological development is highly buffered and that long-lasting consequences of adversity usually are associated with either organic damage or severe interference in the normative protective processes embedded in the

caregiving system. Children who experience chronic adversity fare better or recover more successfully when they have a positive relationship with a competent adult, they are good learners and problem-solvers, they are engaging to other people, and they have areas of competence and perceived efficacy valued by self or society. Future studies of resilience will need to focus on processes that facilitate adaptation. Such studies have the potential to illuminate the range and self-righting properties of, constraints on, and linkages among different aspects of cognitive, emotional, and social development.